

Datasheet for ABIN679508  
**anti-HBV PreS1,PreS2 antibody**[Go to Product page](#)

## 1 Image

## Overview

Quantity:	100 µL
Target:	HBV PreS1,PreS2
Reactivity:	Hepatitis B Virus (HBV)
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HBV PreS1,PreS2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from HBSAG_HBVD2 sub. ayw pre S1/S2 protein
Isotype:	IgG
Cross-Reactivity:	Virus
Cross-Reactivity (Details):	Hepatitis B virus ay/ad
Purification:	Purified by Protein A.

## Target Details

Target:	HBV PreS1,PreS2
Abstract:	<a href="#">HBV PreS1,PreS2 Products</a>
Target Type:	Viral Protein
Background:	Synonyms: Hepatitis B, HBSAG_HBVD2, HBSAG_HBVC5, Large envelope protein, Large surface

## Target Details

protein, L glycoprotein, Major surface antigen, Large S protein, L-HBsAg, LHB

Background: Hepatitis B Virus (HBV) infection induces a disease state characterised by liver damage, inflammation and viral persistence. Infection also increases the risk of hepatocellular carcinoma. HBV belongs to the Hepadnaviridae family of viruses. Its genome consists of partially double stranded circular DNA. The DNA is enclosed in a nucleocapsid, or core antigen (HBcAg), which is surrounded by a spherical envelope (surface antigen or HBsAg). The core antigen shares its sequences with the e antigen (HBeAg) but no cross reactivity between the two proteins has been observed. The HBV genome also encodes a DNA polymerase that also acts as a reverse transcriptase. Hepatitis B infection is normally diagnosed from serological tests that detect HBsAg but as the disease progresses this antigen may no longer be present in the blood and tests for HBcAg are used. If HBsAg can be detected in the blood for longer than six months, chronic hepatitis B is diagnosed. The antigenic determinant of the protein moiety of the HBsAg determines specific characteristics of different serotypes and provides the basis of immunodetection. HBsAg has antigenic heterogeneity, specifically, two pairs of sub specific determinants, d/y and w/r allow the following combinations: adw, ayw, adr, ayr.

## Application Details

Application Notes: WB 1:100-1000  
IHC-P 1:100-500  
IF(IHC-P) 1:50-200

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: 0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: 4 °C,-20 °C

Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Expiry Date: 12 months



**ELISA**

**Image 1.** Antigen: 0.2ug/100ul Primary: Antiserum, 1:500, 1:1000, 1:2000, 1:4000, 1:8000, 1:16000, 1:32000, Secondary: HRP conjugated Goat Anti-Rabbit IgG -HRP) at 1:5000, TMB staining, Read the data in MicroplateReader by 450nm.