

Datasheet for ABIN680617 anti-CACNA1G antibody (AA 901-1000) (HRP)



Go to Product page

_				
()	1//	rv	IO	Λ/
()	VC	. I V	1	v v

Quantity:	100 μL	
Target:	CACNA1G	
Binding Specificity:	AA 901-1000	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CACNA1G antibody is conjugated to HRP	
Application:	ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffinembedded Sections) (IHC (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CACNA1G/Cav31
Isotype:	IgG
Specificity:	Possible cross-reactivity with CACNA1H and CACNA1I
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Pig,Guinea Pig
Purification:	Purified by Protein A.

Target Details

Target: CACNA1G

Target Details

Alternative Name:	CACNA1G (CACNA1G Products)	
Background:	Synonyms: NBR13, Cav3.1, Ca(V)T.1, Voltage-dependent T-type calcium channel subunit alpha-	
	1G, Cav3.1c, Voltage-gated calcium channel subunit alpha Cav3.1, CACNA1G, KIAA1123	
	Background: Voltage-sensitive calcium channels (VSCC) mediate the entry of calcium ions into	
	excitable cells and are also involved in a variety of calcium-dependent processes, including	
	muscle contraction, hormone or neurotransmitter release, gene expression, cell motility, cell	
	division and cell death. The isoform alpha-1G gives rise to T-type calcium currents. T-type	
	calcium channels belong to the "low-voltage activated (LVA)" group and are strongly blocked by	
	mibefradil. A particularity of this type of channel is an opening at quite negative potentials and a	
	voltage-dependent inactivation. T-type channels serve pacemaking functions in both central	
	neurons and cardiac nodal cells and support calcium signaling in secretory cells and vascular	
	smooth muscle. They may also be involved in the modulation of firing patterns of neurons	
	which is important for information processing as well as in cell growth processes.	
Gene ID:	8913	
UniProt:	043497	
Application Details		
Application Notes:	IHC-P 1:200-400	
	IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish	
	peroxidase.	

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months