

Datasheet for ABIN6809888

CSF1R Protein (AA 20-510) (His tag)

2 Images



Go to Product page

(11 /	\sim	× 1	i.	\sim		
\cup	٧	Н	rv	I	H	٧	1

Quantity:	100 μg
Target:	CSF1R
Protein Characteristics:	AA 20-510
Origin:	Dog
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This CSF1R protein is labelled with His tag.

Product Details

Sequence:	AA 20-510
Purity:	>95 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per μg by the LAL method.

Target Details

Target:	CSF1R
Alternative Name:	M-CSF R (CSF1R Products)
Background:	Colony stimulating factor 1 receptor (CSF1R) is also known as macrophage colony-stimulating
	factor receptor (M-CSFR), CD115 Cluster of Differentiation 115 (CD115), C-FMS, CSFR, FIM2,
	FMS, and is a member of the typeIII subfamily of receptor tyrosine kinases (RTKs). CSF1R is a
	receptor for a cytokine called colony stimulating factor 1, The protein encoded by the CSFR1

gene is the receptor for colony stimulating factor 1, a cytokine which controls the production, differentiation, and function of macrophages. This receptor mediates most, if not all, of the biological effects of this cytokine. Ligand binding activates CSFR1 through a process of oligomerization and transphosphorylation . Mutations in CSF1R are associated with chronic myelomonocytic leukemia and type M4 acute myeloblastic leukemia. Increased levels of CSF1R1 are found in microglia in Alzheimer's disease and after brain injuries. The increased receptor expression causes microglia to become more active. Both CSF1R, and its ligand colony stimulating factor 1 play an important role in the development of the mammary gland and may be involved in the process of mammary gland carcinogenesis.

Molecular Weight: 56.1 kDa

NCBI Accession: XP_546306


Pathways: RTK Signaling, Inositol Metabolic Process, Cell-Cell Junction Organization

Application Details

Restrictions: For Research Use only

Handling

Format:	Lyophilized
Buffer:	PBS, pH 7.4
Handling Advice:	Please avoid repeated freeze-thaw cycles.
Storage:	-20 °C



kDa M R 116.0 66.2 45.0 35.0 25.0 18.4

ELISA

Image 1. Immobilized Canine M-CSF R, His Tag (ABIN6731307,ABIN6809888) at 2 μ g/mL (100 μ L/well) can bind Biotinylated Human M-CSF, His Tag, Avi Tag (ABIN6386447,ABIN6388276) with a linear range of 1-16 ng/mL (QC tested).

SDS-PAGE

Image 2. Canine M-CSF R, His Tag on under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95 %.