

Datasheet for ABIN6810503

Histone 3 ELISA Kit

[Go to Product page](#)

Overview

Quantity:	96 tests
Target:	Histone 3 (H3)
Binding Specificity:	H3K4me3
Reactivity:	Human, Mouse, Rat
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details

Purpose:	Human, Mouse and Rat Methylated-Histone H3 (H3K4me3) and Histone H3 ELISA Kit. This ELISA is for measuring Methylated Histone H3 (H3K4me3) and Histone H3 in Human, Mouse and Rat Cell Lysates.
Sample Type:	Cell Lysate, Tissue Lysate
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	The antibody pair provided in this kit recognizes human, mouse and rat Methylated-Histone H3 (H3K4me3).
Characteristics:	• Rapidly measure hydroxylated protein in lysates • Screen numerous different cell lysates without performing a Western Blot analysis • Minimal hands-on time, convenient, and non-radioactive material
Components:	• Pre-Coated 96-well Strip Microplate • Wash Buffer

Product Details

- Anti-hydroxylated Antibody
- HRP-Conjugated Secondary Antibody
- Assay Diluent
- TMB One-Step Substrate
- Stop Solution
- Lysis Buffer
- Positive Control Sample

Material not included:	• Distilled or deionized water • 100 mL and 1 liter graduated cylinders • Tubes to prepare sample dilutions • Protease and Phosphatase inhibitors • Precision pipettes to deliver 2 µL to 1 mL volumes • Adjustable 1-25 mL pipettes for reagent preparation • Benchtop rocker or shaker • Microplate reader capable of measuring absorbance at 450 nm
------------------------	--

Target Details

Target:	Histone 3 (H3)
Alternative Name:	Histone H3

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Protocol:	1. Prepare all reagents and samples as instructed in the manual. 2. Add 100 µL of sample or positive control to each well. 3. Incubate 2.5 h at RT or O/N at 4 °C. 4. Add 100 µL of prepared primary antibody to each well. 5. Incubate 1 h at RT. 6. Add 100 µL of prepared 1X HRP-Streptavidin to each well. 7. Incubate 1 h at RT. 8. Add 100 µL of TMB One-Step Substrate Reagent to each well. 9. Incubate 30 min at RT. 10. Add 50 µL of Stop Solution to each well. 11. Read at 450 nm immediately.

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Storage:	-20 °C
Storage Comment:	Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.
Expiry Date:	6 months