



[Go to Product page](#)

Datasheet for ABIN6810516

FRS2 ELISA Kit

Overview

Quantity: 96 tests

Target: FRS2

Binding Specificity: pTyr436

Reactivity: Human, Mouse

Method Type: Sandwich ELISA

Application: ELISA

Product Details

Purpose: Human and Mouse Phospho-FRS2 (Tyr436) ELISA Kit. This assay semi-quantitatively measures FRS2 phosphorylated at Tyrosine-436 in cell lysate samples.

Sample Type: Cell Samples, Tissue Lysate

Analytical Method: Semi-Quantitative

Detection Method: Colorimetric

Specificity: This ELISA kit recognizes Human and Mouse FRS2 phosphorylated at site Tyrosine-436.

Characteristics:

- Pre-Coated 96-well Strip Microplate
- Wash Buffer
- Anti-Phospho Antibody
- HRP-Conjugated Secondary Antibody
- Assay Diluent
- TMB One-Step Substrate
- Stop Solution
- Lysis Buffer

Product Details

- Positive Control Sample

- Components:
- Pre-Coated 96-well Strip Microplate
 - Wash Buffer
 - Anti-Phospho Antibody
 - HRP-Conjugated Secondary Antibody
 - Assay Diluent
 - TMB One-Step Substrate
 - Stop Solution
 - Lysis Buffer
 - Positive Control Sample

- Material not included:
- Distilled or deionized water
 - 100 mL and 1 liter graduated cylinders
 - Tubes to prepare sample dilutions
 - Protease and Phosphatase inhibitors
 - Precision pipettes to deliver 2 μ L to 1 mL volumes
 - Adjustable 1-25 mL pipettes for reagent preparation
 - Benchtop rocker or shaker
 - Microplate reader capable of measuring absorbance at 450 nm

Target Details

Target: FRS2

Alternative Name: FRS2 ([FRS2 Products](#))

Gene ID: 10818

UniProt: [Q8WU20](#), [Q8C180](#)

Pathways: [Fc-epsilon Receptor Signaling Pathway](#), [EGFR Signaling Pathway](#), [Neurotrophin Signaling Pathway](#), [Signaling Events mediated by VEGFR1 and VEGFR2](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

- Protocol:
1. Prepare all reagents and samples as instructed in the manual.
 2. Add 100 μ L of sample or positive control to each well.
 3. Incubate 2.5 h at RT or O/N at 4 $^{\circ}$ C.
 4. Add 100 μ L of prepared primary antibody to each well.
 5. Incubate 1 h at RT.
 6. Add 100 μ L of prepared 1X HRP-Streptavidin to each well.

Application Details

7. Incubate 1 h at RT.
8. Add 100 μ L of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.
10. Add 50 μ L of Stop Solution to each well.
11. Read at 450 nm immediately.

Restrictions: For Research Use only

Handling

Storage: -20 °C

Storage Comment: Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.

Expiry Date: 6 months