antibodies .- online.com







anti-CYP2C9 antibody (Cy5.5)



| () | 1/0 | r\ /1 | 014 | |
|-----|-----|-------|-----|---|
| () | ve | I V I | -v | V |
| | | | | |

| Quantity: | 100 μL |
|--------------|---|
| Target: | CYP2C9 |
| Reactivity: | Human, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This CYP2C9 antibody is conjugated to Cy5.5 |
| Application: | Western Blotting (WB) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human CYP2C9 |
|-------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse |
| Purification: | Purified by Protein A. |

Target Details

| Target: | CYP2C9 | |
|-------------------|---|--|
| Alternative Name: | CYP2C9 (CYP2C9 Products) | |
| Background: | Synonyms: CPC9, CYP2C, CYP2C10, CYPIIC9, P450IIC9, Cytochrome P450 2C9, (R)-limonene 6-monooxygenase, (S)-limonene 7-monooxygenase, | |
| | Cytochrome P-450MP, Cytochrome P450 MP-4, Cytochrome P450 MP-8, Cytochrome P450 PB-1, S-mephenytoin 4-hydroxylase, CYP2C9 | |

Target Details

Background: Cytochromes P450 are a group of heme-thiolate monooxygenases. In liver microsomes, this enzyme is involved in an NADPH-dependent electron transport pathway. It oxidizes a variety of structurally unrelated compounds, including steroids, fatty acids, and xenobiotics. This enzyme contributes to the wide pharmacokinetics variability of the metabolism of drugs such as S-warfarin, diclofenac, phenytoin, tolbutamide and losartan.

Gene ID:

1559

UniProt:

P11712

Application Details

Application Notes:

IF(IHC-P)(1:100-500)

Optimal working dilution should be determined by the investigator.

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |