antibodies -online.com





anti-RECK antibody (AA 801-900) (Cy5)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	RECK	
Binding Specificity:	AA 801-900	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This RECK antibody is conjugated to Cy5	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human RECK	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Pig,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	RECK	
Alternative Name:	RECK (RECK Products)	
Background:	Synonyms: ST15, Reversion-inducing cysteine-rich protein with Kazal motifs, hRECK,	

Expiry Date:

12 months

l arget Details	
	Suppressor of tumorigenicity 15 protein, RECK
	Background: Negatively regulates matrix metalloproteinase-9 (MMP-9) by suppressing MMP-9
	secretion and by direct inhibition of its enzymatic activity. RECK down-regulation by oncogenic
	signals may facilitate tumor invasion and metastasis. Appears to also regulate MMP-2 and
	MT1-MMP, which are involved in cancer progression.
Gene ID:	8434
UniProt:	095980
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.