antibodies -online.com





anti-RECK antibody (AA 801-900) (Cy7)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	RECK
Binding Specificity:	AA 801-900
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RECK antibody is conjugated to Cy7
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human RECK
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Pig,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	RECK
Alternative Name:	RECK (RECK Products)
Background:	Synonyms: ST15, Reversion-inducing cysteine-rich protein with Kazal motifs, hRECK,

Expiry Date:

12 months

Target Details	
	Suppressor of tumorigenicity 15 protein, RECK
	Background: Negatively regulates matrix metalloproteinase-9 (MMP-9) by suppressing MMP-9
	secretion and by direct inhibition of its enzymatic activity. RECK down-regulation by oncogenic
	signals may facilitate tumor invasion and metastasis. Appears to also regulate MMP-2 and
	MT1-MMP, which are involved in cancer progression.
Gene ID:	8434
UniProt:	095980
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.