antibodies

Datasheet for ABIN681853 anti-Frizzled5/8 antibody (AA 51-150)

4 Images



Overview

Quantity:	100 μL
Target:	Frizzled5/8 (HYFZD5/8)
Binding Specificity:	AA 51-150
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Frizzled5/8 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Frizzled 5
Isotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Pig,Rabbit
Purification:	Purified by Protein A.
Target Details	

Target:

Frizzled5/8 (HYFZD5/8)


Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/4 | Product datasheet for ABIN681853 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

Alternative Name:	Frizzled 5/8 (HYFZD5/8 Products)
Background:	Synonyms: C2orf31, Frizzled homolog 5, Frizzled-5, Frizzled5, Fz 5, Fz-5, Fz5, FZD 5, hFz5,
	FZD5, FZD5_HUMAN, FzE 5, FzE5, HFZ 5, Seven transmembrane receptor frizzled 5, Wnt
	receptor, hFz8, Fz-8, Frizzled-8, FZD8.
	Background: Members of the 'frizzled' gene family encode 7-transmembrane domain proteins
	that are receptors for Wnt signaling proteins. The Frizzled 5 protein is believed to be the
	receptor for the Wnt5A ligand. Frizzled 5 has been reported to be expressed in fetal kidney, feta
	and adult liver, fetal lung, and adult pancreas. ESTs have been isolated from bone, liver/spleen,
	placenta, and prostate libraries. Frizzled 5 was cloned from a retina cDNA library. Receptor for
	Wnt proteins. Component of the Wnt-Fzd-LRP5-LRP6 complex that triggers beta-catenin
	signaling through inducing aggregation of receptor-ligand complexes into ribosome-sized
	signalosomes. The beta-catenin canonical signaling pathway leads to the activation of
	disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and
	activation of Wnt target genes. A second signaling pathway involving PKC and calcium fluxes
	has been seen for some family members, but it is not yet clear if it represents a distinct
	pathway or if it can be integrated in the canonical pathway, as PKC seems to be required for
	Wnt-mediated inactivation of GSK-3 kinase. Both pathways seem to involve interactions with G $\!$
	proteins. May be involved in transduction and intercellular transmission of polarity information
	during tissue morphogenesis and/or in differentiated tissues. Coreceptor along with RYK of
	Wnt proteins, such as WNT1.
Gene ID:	7855, 8325
Pathways:	WNT Signaling
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	FCM 1:20-100
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/4 | Product datasheet for ABIN681853 | 03/07/2024 | Copyright antibodies-online. All rights reserved. Handling


Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months

Images



Western Blotting


Image 1. K562 cell lysates probed with Rabbit Anti-Frizzled 5/8 Polyclonal Antibody, Unconjugated at 1:500 for 90 min at 37°C.



Immunohistochemistry (Paraffin-embedded Sections)

Image 2. Formalin-fixed and paraffin embedded mouse embryo labeled with Anti-Frizzled 5/8 Polyclonal Antibody, Unconjugated at 1:200 followed by conjugation to the secondary antibody and DAB staining.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/4 | Product datasheet for ABIN681853 | 03/07/2024 | Copyright antibodies-online. All rights reserved. Images



SDS-PAGE

Image 3. L1 rat lung lysates, L2 rat brain lysates probed (ABIN681853) Anti-Frizzled-5 Polyclonal, Unconjugated at 1:200 in 4 °C. Followed by conjugation to secondary antibody at 1:3000 90min in 37 °C. Predicted band and observed band size: 62kD.

Please check the product details page for more images. Overall 4 images are available for ABIN681853.