

Datasheet for ABIN682207

anti-IKZF3 antibody (AA 251-350) (HRP)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	IKZF3
Binding Specificity:	AA 251-350
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IKZF3 antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human IKZF3
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	IKZF3
Alternative Name:	IKZF3 (IKZF3 Products)

Target Details

Background:	Synonyms: Aiolos, IKAROS family zinc finger 3 Aiolos, IKAROS family zinc finger 3, Ikaros family zinc finger protein 3, IKZF 3, IKZF3, IKZF3_HUMAN, Zinc finger protein Aiolos, Zinc finger protein subfamily 1A 3 Aiolos, Zinc finger protein subfamily 1A 3, ZNFN1A3. Background: IKZF3 belongs to the Ikaros family of zinc-finger proteins. Members of this family are transcription factors that play an important role in the regulation of lymphocyte differentiation. They are involved in the control of gene expression and when associated with nuclear complexes, participate in nucleosome remodeling. IKZF3 is important in the regulation of B lymphocyte proliferation and differentiation.
-------------	---

Gene ID:	22806
----------	-------

Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months