.-online.com antibodies

Datasheet for ABIN682320 anti-Syncytin A (SYNA) (AA 451-538) antibody (Biotin)



Overview

Quantity:	100 µL
Target:	Syncytin A (SYNA)
Binding Specificity:	AA 451-538
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	Biotin
Application:	ELISA, Western Blotting (WB)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Syncytin 1	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse	
Purification:	Purified by Protein A.	

Target Details

Target:	Syncytin A (SYNA)	
Alternative Name:	Syncytin 1 (SYNA Products)	
Background:	Synonyms: ENV, ENVW, HERVW, ERVWE1, HERV7Q, HERV-7q, HERVWENV, HERV-W-ENV,	
	Syncytin-1, Endogenous retrovirus group W member 1, Env-W, Envelope polyprotein gPr73,	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN682320 | 03/06/2024 | Copyright antibodies-online. All rights reserved.

Enverin, HERV-7q Envelope protein, HERV-W envelope protein, HERV-W_7q21.2 provirus
ancestral Env polyprotein, Syncytin, ERVW-1

Background: This endogenous retroviral envelope protein has retained its original fusogenic properties and participates in trophoblast fusion and the formation of a syncytium during placenta morphogenesis. May induce fusion through binding of SLC1A4 and SLC1A5 (PubMed:10708449, PubMed:12050356, PubMed:23492904). Endogenous envelope proteins may have kept, lost or modified their original function during evolution. Retroviral envelope proteins mediate receptor recognition and membrane fusion during early infection. The surface protein (SU) mediates receptor recognition, while the transmembrane protein (TM) acts as a class I viral fusion protein. The protein may have at least 3 conformational states: pre-fusion native state, pre-hairpin intermediate state, and post-fusion hairpin state. During viral and target cell membrane fusion, the coiled coil regions (heptad repeats) assume a trimer-of-hairpins structure, positioning the fusion peptide in close proximity to the C-terminal region of the ectodomain. The formation of this structure appears to drive apposition and subsequent fusion of membranes.

Gene ID:	30816
UniProt:	Q9UQF0

Application Details

Application Notes:	WB 1:300-5000	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C for 12 months.	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN682320 | 03/06/2024 | Copyright antibodies-online. All rights reserved.

1.1	(1:
Н	land	ling
		3

Expiry Date:

12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN682320 | 03/06/2024 | Copyright antibodies-online. All rights reserved.