antibodies -online.com





anti-CITED1 antibody (AA 121-193) (Biotin)



Go to Product pag

| () | 11/0 | K\ / | iew | 1 |
|-----|--------|--------|---------------------|---|
| | \cup | 'I V/I | $\square \vee \vee$ | ı |
| | | | | |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | CITED1 |
| Binding Specificity: | AA 121-193 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This CITED1 antibody is conjugated to Biotin |
| Application: | Western Blotting (WB) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human CITED1 | |
|-------------------|--|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Human, Mouse | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | CITED1 |
|--|--|
| Alternative Name: | CITED1 (CITED1 Products) |
| Background: Synonyms: MSG1, Cbp/p300-interacting transactivator 1, Melanocyte-specific protein | |
| | Background: Transcriptional coactivator of the p300/CBP-mediated trancription complex. |

Enhances SMAD-mediated transcription by strengthening the functional link between the DNA-binding SMAD transcription factors and the p300/CBP transcription coactivator complex. Stimulates estrogen-dependent transactivation activity mediated by estrogen receptors signaling, stabilizes the interaction of estrogen receptor ESR1 and histone acetyltransferase EP300. Positively regulates TGF-beta signaling through its association with the SMAD/p300/CBP-mediated transcriptional coactivator complex. Induces transcription from estrogen-responsive promoters and protection against cell death. Potentiates EGR2-mediated transcriptional activation activity from the ERBB2 promoter. Acts as an inhibitor of osteoblastic mineralization through a cAMP-dependent parathyroid hormone receptor signaling. May play a role in pigmentation of melanocytes. Associates with chromatin to the estrogen-responsive TGF-alpha promoter region in a estrogen-dependent manner.

Gene ID:

4435

UniProt:

Q99966

Application Details

Application Notes:

WB 1:300-5000

IHC-P 1:200-400

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C for 12 months. |
| Expiry Date: | 12 months |