# antibodies -online.com





## anti-MAP3K11 antibody (AA 355-460) (Biotin)



Go to Product page

( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL
Target:	MAP3K11
Binding Specificity:	AA 355-460
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MAP3K11 antibody is conjugated to Biotin
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human MLK3
Isotype:	IgG
Specificity:	Due to the similarity of this protein to MLK1 2 and 4, there is a chance that this antibody will cross-react based on homology.
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Cow,Horse
Purification:	Purified by Protein A.

### **Target Details**

Target:	MAP3K11	
Alternative Name:	MLK3 (MAP3K11 Products)	
Background:	Synonyms: MLK3, PTK1, SPRK, MLK-3, MEKK11, Mitogen-activated protein kinase kinase	
	kinase 11, Mixed lineage kinase 3, Src-homology 3 domain-containing proline-rich kinase,	
	MAP3K11	
	Background: Activates the JUN N-terminal pathway. Required for serum-stimulated cell	
	proliferation and for mitogen and cytokine activation of MAPK14 (p38), MAPK3 (ERK) and	
	MAPK8 (JNK1) through phosphorylation and activation of MAP2K4/MKK4 and MAP2K7/MKK7.	
	Plays a role in mitogen-stimulated phosphorylation and activation of BRAF, but does not	
	phosphorylate BRAF directly. Influences microtubule organization during the cell cycle.	
Molecular Weight:	94kDa	
Gene ID:	4296	
UniProt:	Q16584	
Pathways:	MAPK Signaling, Interferon-gamma Pathway	
Application Details		
Application Notes:	WB 1:300-5000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	

## Handling

Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months