

Datasheet for ABIN682903  
**anti-BRCA1 antibody (pSer1524)**



[Go to Product page](#)

2 Images

## Overview

Quantity:	100 µL
Target:	BRCA1
Binding Specificity:	pSer1524
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BRCA1 antibody is un-conjugated
Application:	ELISA, Flow Cytometry (FACS)

## Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human BRCA1 around the phosphorylation site of Ser1524
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified by Protein A.

## Target Details

Target:	BRCA1
Alternative Name:	BRCA1 ( <a href="#">BRCA1 Products</a> )
Background:	Synonyms: IRIS, PSCP, BRCAI, BRCC1, FANCS, PNCA4, RNF53, BROVCA1, PPP1R53, Breast

## Target Details

---

cancer type 1 susceptibility protein, RING finger protein 53, BRCA1

Background: E3 ubiquitin-protein ligase that specifically mediates the formation of 'Lys-6'-linked polyubiquitin chains and plays a central role in DNA repair by facilitating cellular responses to DNA damage. It is unclear whether it also mediates the formation of other types of polyubiquitin chains. The E3 ubiquitin-protein ligase activity is required for its tumor suppressor function. The BRCA1-BARD1 heterodimer coordinates a diverse range of cellular pathways such as DNA damage repair, ubiquitination and transcriptional regulation to maintain genomic stability. Regulates centrosomal microtubule nucleation. Required for normal cell cycle progression from G2 to mitosis. Required for appropriate cell cycle arrests after ionizing irradiation in both the S-phase and the G2 phase of the cell cycle. Involved in transcriptional regulation of P21 in response to DNA damage. Required for FANCD2 targeting to sites of DNA damage. May function as a transcriptional regulator. Inhibits lipid synthesis by binding to inactive phosphorylated ACACA and preventing its dephosphorylation. Contributes to homologous recombination repair (HRR) via its direct interaction with PALB2, fine-tunes recombinational repair partly through its modulatory role in the PALB2-dependent loading of BRCA2-RAD51 repair machinery at DNA breaks. Component of the BRCA1-RBBP8 complex which regulates CHEK1 activation and controls cell cycle G2/M checkpoints on DNA damage via BRCA1-mediated ubiquitination of RBBP8. Acts as a transcriptional activator (PubMed:20160719).

---

Gene ID: 672

---

UniProt: [P38398](#)

---

Pathways: [Cell Division Cycle, DNA Damage Repair, Intracellular Steroid Hormone Receptor Signaling Pathway, Positive Regulation of Response to DNA Damage Stimulus](#)

## Application Details

---

Application Notes: WB 1:300-5000  
ELISA 1:500-1000  
FCM 1:20-100

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

---


Concentration: 1 µg/µL

---

## Handling


Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months

## Images



### Immunohistochemistry

**Image 1.** Formalin-fixed and paraffin embedded human colon carcinoma labeled with Rabbit Anti Phospho-BRCA1(Ser1524) Polyclonal Antibody, Unconjugated (ABIN682903) at 1:200 followed by conjugation to the secondary antibody and DAB staining



**Image 2.** HL-60 cells were fixed with 4% PFA for 10min at room temperature, 0.1% PBST for 20 min at room temperature, and incubated in 5% BSA blocking buffer for 30 min at room temperature. Cells were then stained with BRCA1 (Ser1524) Polyclonal Antibody at 1: 50 dilution in blocking buffer and incubated for 30 min at room temperature, washed twice with 2%BSA in PBS, followed by secondary antibody incubation for 40 min at room temperature. Acquisitions of 20, 000 events were performed.