

Datasheet for ABIN684313

anti-Doublecortin antibody (pSer47)



Overview

Quantity:	100 μL
Target:	Doublecortin (DCX)
Binding Specificity:	pSer47
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Doublecortin antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Flow Cytometry (FACS), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunofluorescence (Cultured Cells) (IF (cc))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human Doublecortin around the phosphorylation site of Ser47
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	Doublecortin (DCX)
Alternative Name:	Doublecortin (DCX Products)
Background:	Synonyms: Doublecortex, DBCN, Dbct, DC, Doublin, Lis X, Lissencephalin X, Lissencephaly X linked, Lissencephaly X linked doublecortin, LISX, Neuronal migration protein doublecortin, SCLH, XLIS, p-DCX, p-Doublecortex, phospho-Doublecortex. Background: Neuronal Marker Doublecortin (DCX) is a microtubule-associated protein expressed almost exclusively in immature neurons. Neuronal precursors begin to express DCX shortly after exiting the cell cycle, and continue to express DCX for 2-3 weeks as the cells mature into neurons. Downregulation of DCX begins after 2 weeks, and occurs at the same time that these cells begin to express, a marker for mature neurons. Due to the nearly exclusive expression of DCX in developing neurons, this protein has been used increasingly as a marker for neurogenesis. Indeed, the levels of DCX expression increase in response to exercise, which occurs in parallel with increased BrdU labelling, currently a "gold standard" in measuring neurogenesis.
Gene ID:	1641
UniProt:	043602

Application Details

Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	FCM 1:20-100
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months