

Datasheet for ABIN685053

anti-FAK antibody (pTyr576, pTyr577) (Cy5.5)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	FAK (PTK2)
Binding Specificity:	pTyr576, pTyr577
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FAK antibody is conjugated to Cy5.5
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Flow Cytometry (FACS)

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human FAK around the phosphorylation site of Tyr576/577
Isotype:	IgG
Specificity:	These phosphorylation sites are homologous to Tyr614 + Tyr615 in Mouse.
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	FAK (PTK2)
Alternative Name:	FAK + (PTK2 Products)
Background:	Synonyms: FAK, FADK, FAK1, FRNK, PPP1R71, p125FAK, pp125FAK, Focal adhesion kinase 1, FADK 1, Focal adhesion kinase-related nonkinase, Protein phosphatase 1 regulatory subunit 71, Protein-tyrosine kinase 2, PTK2 Background: Non-receptor protein-tyrosine kinase that plays an essential role in regulating cell migration, adhesion, spreading, reorganization of the actin cytoskeleton, formation and disassembly of focal adhesions and cell protrusions, cell cycle progression, cell proliferation and apoptosis. Required for early embryonic development and placenta development. Required for embryonic angiogenesis, normal cardiomyocyte migration and proliferation, and normal heart development. Regulates axon growth and neuronal cell migration, axon branching and synapse formation, required for normal development of the nervous system. Plays a role in osteogenesis and differentiation of osteoblasts. Functions in integrin signal transduction, but also in signaling downstream of numerous growth factor receptors, G-protein coupled receptors (GPCR), EPHA2, netrin receptors and LDL receptors. Forms multisubunit signaling complexes with SRC and SRC family members upon activation, this leads to the phosphorylation of additional tyrosine residues, creating binding sites for scaffold proteins, effectors and substrates. Regulates numerous signaling pathways. Promotes activation of phosphatidylinositol 3-kinase and the AKT1 signaling cascade. Promotes activation of MAPK1/ERK2, MAPK3/ERK1 and the MAP kinase signaling cascade. Promotes localized and transient activation of guanine nucleotide exchange factors (GEFs) and GTPase-activating proteins (GAPs), and thereby modulates the activity of Rho family GTPases. Signaling via CAS family members mediates activation of RAC1. Recruits the ubiquitin ligase MDM2 to P53/TP53 in the nucleus, and thereby regulates P53/TP53 activity, P53/TP53 ubiquitination and proteasomal degradation.
Gene ID:	5747
UniProt:	Q05397
Pathways:	Response to Growth Hormone Stimulus , CXCR4-mediated Signaling Events , Smooth Muscle Cell Migration , Signaling of Hepatocyte Growth Factor Receptor , VEGF Signaling

Application Details

Application Notes:	FCM 1:20-100 IF(IHC-P) 1:50-200
--------------------	------------------------------------

Application Details

	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months