

Datasheet for ABIN685235

anti-HDAC3 antibody (pSer424) (FITC)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	HDAC3
Binding Specificity:	pSer424
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HDAC3 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human HDAC3 around the phosphorylation site of Ser424
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

Target Details

Target:	HDAC3
Alternative Name:	HDAC3 (HDAC3 Products)

Target Details

Background:	Synonyms: HDAC3 phospho S424, HDAC3 phospho Ser424, p-HDAC3 Ser424, RPD3 2, RPD3-2, HD 3, HD3, HDAC 3, Histone deacetylase 3 HD3 RPD3-2, histone deacetylase 3, RPD 3, RPD3, SMAP 45, SMAP-45, SMAP45, HDAC3_HUMAN. Background: Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to the histone deacetylase/acuc/apha family. It has histone deacetylase activity and represses transcription when tethered to a promoter. It may participate in the regulation of transcription through its binding with the zinc-finger transcription factor YY1. This protein can also down-regulate p53 function and thus modulate cell growth and apoptosis. This gene is regarded as a potential tumor suppressor gene. [provided by RefSeq, Jul 2008].
Gene ID:	8841
Pathways:	Neurotrophin Signaling Pathway , Regulation of Lipid Metabolism by PPARalpha , Regulation of Muscle Cell Differentiation , Skeletal Muscle Fiber Development

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months