

Datasheet for ABIN685363

anti-TANK antibody (AA 151-260) (Cy7)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	TANK
Binding Specificity:	AA 151-260
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TANK antibody is conjugated to Cy7
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human TANK
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat
Purification:	Purified by Protein A.

Target Details

Target:	TANK
Alternative Name:	TANK (TANK Products)
Background:	Synonyms: I TRAF, ITRAF, TRAF family member associated NF KAPPA B activator, TRAF family

Target Details

member associated NFKB activator, TRAF interacting protein, TRAF interacting protein TANK isoform a, I-TRAF, Tank, TANK_HUMAN, TRAF family member-associated NF-kappa-B activator, TRAF-interacting protein, TRAF interacting protein TANK isoform b, TRAF2.

Background: TANK was initially identified as a novel TRAF-interacting protein that regulated TRAF-mediated signal transduction. Specifically, ligand binding by surface receptors in the tumor necrosis factor (TNF) receptor and Toll/interleukin-1 (IL-1) receptor families lead to the formation of a TRAF/TANK complex that mediates the activation of the transcription factor NF-kappaB. TANK is found in the cytoplasm and can bind to TRAF1, TRAF2, or TRAF3, thereby inhibiting TRAF function by sequestering the TRAFs in a latent state in the cytoplasm. For example, this protein can block TRAF2 binding to LMP1, the Epstein Barr virus transforming protein, and inhibit LMP1-mediated NF kappa B activation.

Gene ID: 10010

Pathways: [p53 Signaling](#), [TLR Signaling](#), [Activation of Innate immune Response](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months