

Datasheet for ABIN685914

anti-COPS5 antibody (AA 221-334) (Biotin)



Go to Product page

_					
	W	0	rv	10	W

Quantity:	100 μL
Target:	COPS5
Binding Specificity:	AA 221-334
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This COPS5 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)),
	Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Product Details Immunogen:	KLH conjugated synthetic peptide derived from human Jab1
	KLH conjugated synthetic peptide derived from human Jab1 IgG
lmmunogen:	
Immunogen: Isotype:	IgG
Immunogen: Isotype: Cross-Reactivity:	IgG Human
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity:	IgG Human Mouse,Rat,Dog,Cow,Pig,Horse
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity: Purification:	IgG Human Mouse,Rat,Dog,Cow,Pig,Horse

Target Details

Background:	Synonyms: 38 kDa Mov34 homolog, COP9 constitutive photomorphogenic homolog subunit 5, COP9 constitutive photomorphogenic homolog subunit 5, COP9 signalosome complex subunit 5, COP9 signalosome subunit 5, COP9 signalosome subunit 5, COPS 5, COPS5, CSN 5, CSN5, JAB 1, Jun activation domain binding protein 1, Jun activation domain-binding protein, MGC3149, MOV 34, MOV34, SGN 5, SGN5, Signalosome subunit 5, CSN5_HUMAN. Background: The protein encoded by this gene is one of the eight subunits of COP9 signalosome, a highly conserved protein complex that functions as an important regulator in multiple signaling pathways. The structure and function of COP9 signalosome is similar to that of the 19S regulatory particle of 26S proteasome. COP9 signalosome has been shown to interact with SCF-type E3 ubiquitin ligases and act as a positive regulator of E3 ubiquitin ligases. This protein is reported to be involved in the degradation of cyclin-dependent kinase inhibitor CDKN1B/p27Kip1. It is also known to be an coactivator that increases the specificity of
Gene ID:	JUN/AP1 transcription factors. 10987
Pathways:	Cell Division Cycle
Application Details	
Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.

		1.	
\vdash	land	ling	
1 1	ıaııu	11111	۰

Expiry Date:

12 months