



[Go to Product page](#)

Datasheet for ABIN687460

## anti-CXCL14 antibody (AA 51-99) (Cy3)

### Overview

|                      |   |
|----------------------|---|
| Quantity:            | 100 µL  |
| Target:              | CXCL14  |
| Binding Specificity: | AA 51-99  |
| Reactivity:          | Human, Rat  |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This CXCL14 antibody is conjugated to Cy3   |
| Application:         | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

### Product Details

|                       |  |
|-----------------------|--|
| Immunogen:            | KLH conjugated synthetic peptide derived from rat CXCL14 |
| Isotype:              | IgG  |
| Cross-Reactivity:     | Human, Rat   |
| Predicted Reactivity: | Mouse  |
| Purification:         | Purified by Protein A.                                   |

### Target Details

|                   |  |
|-------------------|--|
| Target:           | CXCL14                                     |
| Alternative Name: | CXCL14 ( <a href="#">CXCL14 Products</a> ) |

## Target Details

---

**Background:** Synonyms: C-X-C mot chemokine 14, Chemokine BRAK, MIP-2G, Small-inducible cytokine B14  
Background: This gene belongs to the cytokine gene family which encode secreted proteins involved in immunoregulatory and inflammatory processes. The protein encoded by this gene is structurally related to the CXC (Cys-X-Cys) subfamily of cytokines. Members of this subfamily are characterized by two cysteines separated by a single amino acid. This cytokine displays chemotactic activity for monocytes but not for lymphocytes, dendritic cells, neutrophils or macrophages. It has been implicated that this cytokine is involved in the homeostasis of monocyte-derived macrophages rather than in inflammation. [provided by RefSeq]. [FUNCTION] Potent chemoattractant for neutrophils, and weaker for dendritic cells. Not chemotactic for T-cells, B-cells, monocytes, natural killer cells or granulocytes. Does not inhibit proliferation of myeloid progenitors in colony formation assays.

**Gene ID:** 306748

**Pathways:** [Autophagy](#)

## Application Details

---

**Application Notes:** IF(IHC-P) 1:50-200  
IF(IHC-F) 1:50-200  
IF(ICC) 1:50-200

**Restrictions:** For Research Use only

## Handling

---

**Format:** Liquid

**Concentration:** 1 µg/µL

**Buffer:** Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

**Preservative:** ProClin

**Precaution of Use:** This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

**Storage:** -20 °C

**Storage Comment:** Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

**Expiry Date:** 12 months