

Datasheet for ABIN6936330

anti-ABCC1 antibody



Overview

Quantity:	100 μg
Target:	ABCC1
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This ABCC1 antibody is un-conjugated
Application:	Western Blotting (WB), Flow Cytometry (FACS), ELISA, Coating (Coat)

Product Details

Immunogen:	Recombinant full-length human ABCC1 protein
Clone:	MRP1-1343
Isotype:	IgG1 kappa
Specificity:	The two members of the large family of ABC transporters known to confer multidrug resistance in human cancer cells are the MDR1 P-glycoprotein and the multidrug-resistance protein MRP1. MRP1 is an integral membrane protein that contains an MDR-like core, an N-terminal membrane-bound region and a cytoplasmic linker, and it is expressed in various cerebral cells, as well as in lung, testis and peripheral blood.
Cross-Reactivity (Details):	Human.
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target:	ABCC1
Alternative Name:	ABCC1 (ABCC1 Products)
Background:	ABC29, ABCC1, ATP binding cassette sub family C (CFTR/MRP) member 1, ATP-binding
	cassette sub-family C member 1, GSX, Leukotriene C(4) transporter, LTC4 transporter,
	Multidrug resistance-associated protein 1 (MRP1), MRP1 / ABCC1 (Multidrug Resistance
	Related Protein 1)
	Cellular localisation: Cytoplasmic and Cell Surface
Molecular Weight:	190kDa
Gene ID:	4363, 391464
UniProt:	P33527
Pathways:	SARS-CoV-2 Protein Interactome
Application Details	
Application Notes:	Known_Application: ELISA (For coating, order antibody without BSA), Flow Cytometry (1-2 μ
	Known_Application: ELISA (For coating, order antibody without BSA), Flow Cytometry (1-2 μ g/million cells), Western Blot (1-2 μ g/mL), Optimal dilution for a specific application should be
	g/million cells), Western Blot (1-2 μg/mL), Optimal dilution for a specific application should be
	g/million cells), Western Blot (1-2 μ g/mL), Optimal dilution for a specific application should be determined.
Application Notes:	g/million cells), Western Blot (1-2 µg/mL), Optimal dilution for a specific application should be determined. Positive_Control: A549 cells. Colon, Endometrium, Ovary. Testis or Tonsil.
Application Notes: Restrictions:	g/million cells), Western Blot (1-2 µg/mL), Optimal dilution for a specific application should be determined. Positive_Control: A549 cells. Colon, Endometrium, Ovary. Testis or Tonsil.
Application Notes: Restrictions: Handling	g/million cells), Western Blot (1-2 µg/mL), Optimal dilution for a specific application should be determined. Positive_Control: A549 cells. Colon, Endometrium, Ovary. Testis or Tonsil. For Research Use only
Application Notes: Restrictions: Handling Concentration:	g/million cells), Western Blot (1-2 µg/mL), Optimal dilution for a specific application should be determined. Positive_Control: A549 cells. Colon, Endometrium, Ovary. Testis or Tonsil. For Research Use only 1.0 mg/mL
Application Notes: Restrictions: Handling Concentration: Buffer:	g/million cells), Western Blot (1-2 µg/mL), Optimal dilution for a specific application should be determined. Positive_Control: A549 cells. Colon, Endometrium, Ovary. Testis or Tonsil. For Research Use only 1.0 mg/mL Prepared in 10 mM PBS, WITHOUT BSA and Azide.
Application Notes: Restrictions: Handling Concentration: Buffer: Preservative:	g/million cells), Western Blot (1-2 µg/mL), Optimal dilution for a specific application should be determined. Positive_Control: A549 cells. Colon, Endometrium, Ovary. Testis or Tonsil. For Research Use only 1.0 mg/mL Prepared in 10 mM PBS, WITHOUT BSA and Azide. Azide free