

Datasheet for ABIN6936751
anti-CD5 antibody (AA 269-366)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	CD5
Binding Specificity:	AA 269-366
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD5 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (Formalin-fixed Sections) (IHC (f))

Product Details

Immunogen:	Recombinant fragment of human CD5 protein (around aa 269-366) (exact sequence is proprietary)
Clone:	CD5-2416
Isotype:	IgG2b kappa
Specificity:	Recognizes a 67 kDa transmembrane protein, which is identified as CD5. The CD5 antigen is found on 95 % of thymocytes and 72 % of peripheral blood lymphocytes. In lymph nodes, the main reactivity is observed in T cell areas. Anti-CD5 is a pan T-cell marker that also reacts with a range of neoplastic B-cells, e.g. chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL), mantle cell lymphoma, and a subset (~10 %) of diffuse large B-cell lymphoma. CD5 aberrant expression is useful in making a diagnosis of mature T-cell neoplasms. Anti-CD5

Product Details

detection is diagnostic in CLL/SLL within a panel of other B-cell markers, especially one that includes anti-CD23. Anti-CD5 is also very useful in differentiating among mature small lymphoid cell malignancies. In addition, anti-CD5 can be used in distinguishing thymic carcinoma (+) from thymoma (-). Anti-CD5 does not react with granulocytes or monocytes.

Cross-Reactivity (Details): Human.

Purification: 1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target: CD5

Alternative Name: CD5 ([CD5 Products](#))

Background: CD5 antigen (p56 62), LEU1, Ly12, LyA, Lymphocyte antigen T1/Leu-1, Lymphocyte glycoprotein T1/Leu1, T-cell surface glycoprotein CD5,CD5 (Mantle Cell Lymphoma Marker)
Cellular localisation: Cell Surface

Molecular Weight: 67kDa

Gene ID: 921, 58685

UniProt: [P06127](#)

Application Details

Application Notes: Known_Application: Flow Cytometry (1-2 µg/million cells), Western Blot (1-2 µg/mL), Immunofluorescence (1-2 µg/million cells),Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes), Optimal dilution for a specific application should be determined.
Positive_Control: 293T, Ramos or MOLT-4 cells. Tonsil or Lymph Node.

Restrictions: For Research Use only

Handling

Concentration: 1.0 mg/mL

Buffer: Prepared in 10 mM PBS, WITHOUT BSA and Azide.

Preservative: Azide free

Storage: -20 °C,-80 °C

Handling

Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.
Expiry Date:	24 months