

Datasheet for ABIN6936922
anti-CDX2 antibody (AA 150-249)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	CDX2
Binding Specificity:	AA 150-249
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CDX2 antibody is un-conjugated
Application:	ELISA, Coating (Coat), Immunohistochemistry (Formalin-fixed Sections) (IHC (f)), Immunostaining (Ist)

Product Details

Immunogen:	Recombinant fragment (around aa150-249) of human CDX2 protein (exact sequence is proprietary)
Clone:	CDX2-1690
Isotype:	IgG2a kappa
Specificity:	The intestine-specific transcription factors CDX1 and CDX2 are important for directing intestinal development, differentiation, proliferation and maintenance of the intestinal phenotype. CDX2 protein expression has been seen in GI carcinomas. Anti-CDX2 has been useful to establish GI origin of metastatic adenocarcinomas and carcinoids and is especially useful to distinguish metastatic colorectal adenocarcinoma from lung adenocarcinoma. However, mucinous carcinomas of the ovary also express CDX2 protein. It limits the usefulness of this marker in the

Product Details

	distinction of metastatic colorectal adenocarcinoma from mucinous carcinoma of the ovary.
Cross-Reactivity (Details):	Human.
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target:	CDX2
Alternative Name:	CDX2 (CDX2 Products)
Background:	Caudal type homeobox 2, Caudal type homeobox transcription factor 2, Caudal-type homeobox protein 2, CDX2,CDX2 / Caudal Type Homeobox 2 (GI Epithelial Marker) Cellular localisation: Nuclear
Molecular Weight:	40kDa
Gene ID:	1045, 174249
UniProt:	Q99626
Pathways:	Peptide Hormone Metabolism , Stem Cell Maintenance

Application Details

Application Notes:	Known_Application: ELISA (Use Ab at 2-4 µg/mL for coating) (Order Ab without BSA), ,Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 min at RT),(Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0 for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined. Positive_Control: HT29 cells. Colon Carcinoma.
Restrictions:	For Research Use only

Handling

Concentration:	1.0 mg/mL
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.
Preservative:	Azide free
Storage:	-20 °C,-80 °C
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.

Handling

Expiry Date: 24 months