

Datasheet for ABIN6936954
anti-CFTR antibody (AA 258-385)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	CFTR
Binding Specificity:	AA 258-385
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CFTR antibody is un-conjugated
Application:	Immunohistochemistry (Formalin-fixed Sections) (IHC (f)), Flow Cytometry (FACS)

Product Details

Immunogen:	Recombinant fragment (around aa 258-385) of human CFTRprotein (exact sequence is proprietary)
Clone:	CFTR-1643
Isotype:	IgG2b kappa
Specificity:	Recognizes a protein of 165-170kDa, identified as cystic fibrosis transmembrane conductance regulator (CFTR). CFTR is composed of two membrane-spanning domains (MSD), two nucleotide-binding domains (NBD), and an R domain. It is structurally similar to multidrug resistance (Mdr1) protein and both are members of the superfamily of ATP-binding cassette (ABC) transporters, also known as traffic ATPases, which are implicated in the movement of various substrates. The CFTR protein is a small conductance adenosine 3',5'-cyclic monophosphate (cAMP)-activated chloride ion channel found in the apical membranes of

Product Details

epithelia within the pancreas, airway, intestine, bile duct, sweat gland, and male genital ducts.
CFTR is a valuable marker of human pancreatic duct cell development and differentiation.

Cross-Reactivity (Details): Human.

Purification: 1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target: CFTR

Alternative Name: CFTR ([CFTR Products](#))

Background: ABC35, ATP Binding Cassette Superfamily C Member 7 (ABCC7), cAMP-dependent chloride channel, CFTR/MRP, Channel conductance-controlling ATPase, Cystic Fibrosis Transmembrane Conductance Regulator, MRP7, TNR CFTR,CFTR (Cystic Fibrosis Transmembrane Conductance Regulator)
Cellular localisation: Cell surface. Cytoplasm.

Molecular Weight: 165-170kDa

Gene ID: 1080, 489786, 621460

UniProt: [P13569](#)

Application Details

Application Notes: Known_Application: Flow Cytometry (1-2 µg/million cells),Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined.
Positive_Control: MOLT-4 cells. Human pancreas, kidney or placenta.

Restrictions: For Research Use only

Handling

Concentration: 1.0 mg/mL

Buffer: Prepared in 10 mM PBS, WITHOUT BSA and Azide.

Preservative: Azide free

Storage: -20 °C,-80 °C

Handling

Storage Comment: Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.

Expiry Date: 24 months