

## Datasheet for ABIN6936996

# anti-CKB antibody



#### Overview

Quantity:	100 μg
Target:	CKB
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CKB antibody is un-conjugated
Application:	ELISA, Immunofluorescence (IF), Immunohistochemistry (Frozen Sections) (IHC (fro)), Flow Cytometry (FACS), Coating (Coat)

## **Product Details**

Immunogen:

Specificity:  The specificity of this monoclonal antibody to its intended target was validated by HuProtTM  Array, containing more than 19,000, full-length human proteins. The specificity of this  monoclonal antibody to its intended target was validated by HuProt Array containing more than  19,000 full-length, correctly-folded human proteins. Creatine kinases (CK) are a large family of  isoenzymes that regulate levels of ATP in subcellular compartments, where they provide ATP at  sites of fluctuating energy demand by the transfer of phosphates between creatine and adenine nucleotides. CKs provide the energy of phosphate hydrolysis necessary to drive the normal function of many cellular systems. In cells, the cytosolic CK enzymes consist of two subunits,
Array, containing more than 19,000, full-length human proteins. The specificity of this monoclonal antibody to its intended target was validated by HuProt Array containing more than 19,000 full-length, correctly-folded human proteins. Creatine kinases (CK) are a large family of isoenzymes that regulate levels of ATP in subcellular compartments, where they provide ATP at sites of fluctuating energy demand by the transfer of phosphates between creatine and adenine nucleotides. CKs provide the energy of phosphate hydrolysis necessary to drive the normal
monoclonal antibody to its intended target was validated by HuProt Array containing more than 19,000 full-length, correctly-folded human proteins. Creatine kinases (CK) are a large family of isoenzymes that regulate levels of ATP in subcellular compartments, where they provide ATP at sites of fluctuating energy demand by the transfer of phosphates between creatine and adenine nucleotides. CKs provide the energy of phosphate hydrolysis necessary to drive the normal
19,000 full-length, correctly-folded human proteins. Creatine kinases (CK) are a large family of isoenzymes that regulate levels of ATP in subcellular compartments, where they provide ATP at sites of fluctuating energy demand by the transfer of phosphates between creatine and adenine nucleotides. CKs provide the energy of phosphate hydrolysis necessary to drive the normal
isoenzymes that regulate levels of ATP in subcellular compartments, where they provide ATP at sites of fluctuating energy demand by the transfer of phosphates between creatine and adenine nucleotides. CKs provide the energy of phosphate hydrolysis necessary to drive the normal
sites of fluctuating energy demand by the transfer of phosphates between creatine and adenine nucleotides. CKs provide the energy of phosphate hydrolysis necessary to drive the normal
nucleotides. CKs provide the energy of phosphate hydrolysis necessary to drive the normal
function of many callular eyetems. In calls, the cytosolic CK enzymes consist of two subunits
function of many centural systems. In cens, the cytosonic of enzymes consist of two subunits,
which can be either B (brain type) or M (muscle type). There are three different isoenzymes:

Human CKBB protein

### **Product Details**

	CKMM, CKBB and CKMB. This MAb recognizes the CKBB isoenzyme and does not react with	
	the B subunit in CKMB. It shows minimal reactivity with other human serum proteins.	
Cross-Reactivity (Details):	Human. Predicted to react with Chimpanzee, Rhesus Monkey, Dog, Cow, Mouse, Rat, Chicken,	
	Zebrafish, Frog.	
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.	
Target Details		
Target:	CKB	
Alternative Name:	CKB (CKB Products)	
Background:	B-CK, BB-CK, creatine kinase-B, Creatine kinase B chain, CKBB creatine kinase B-type, creatine	
	kinase, brain, CKBB, Brain creatine kinase, Creatine kinase B-type, Creatine Kinase BB	
	Isoenzyme, Creatine kinase brain, Creatine phosphokinase BB,Creatine Kinase-BB (CK-BB)	
	Cellular localisation: Cytoplasmic	
Molecular Weight:	43kDa (Monomer), 86kDa (Dimer)	
Gene ID:	1152, 173724	
UniProt:	P12277	
Application Details		
Application Notes:	Known_Application: ELISA (For coating, order Ab without BSA),Flow Cytometry (1-2 µg/million	
	cells), Immunofluorescence (1-2 μg/mL), Immunohistochemistry (Frozen Only) (1-2 μg/mL for	
	30 minutes at RT) Optimal dilution for a specific application should be determined.	
	Positive_Control: Cerebellum.	
Restrictions:	For Research Use only	
Handling		
Concentration:	1.0 mg/mL	
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.	
Preservative:	Azide free	
Storage:	-20 °C,-80 °C	
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.	

$\vdash$	land	lına
	iaria	шц

Expiry Date:

24 months