

Datasheet for ABIN6937157

Recombinant anti-EPO antibody (AA 28-162)[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	EPO
Binding Specificity:	AA 28-162
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This EPO antibody is un-conjugated
Application:	ELISA, Coating (Coat), Immunohistochemistry (Formalin-fixed Sections) (IHC (f))

Product Details

Immunogen:	Recombinant fragment (around aa 28-162) of human EPO protein (exact sequence is proprietary)
Clone:	EPO-3793R
Isotype:	IgG
Specificity:	Recognizes a protein of about 37 kDa, which is identified as Erythropoietin (EPO). Erythropoietin is a secreted, glycosylated cytokine hormone composed of four alpha helical bundles. It is the primary factor responsible for regulating erythropoiesis during steady-state conditions and in response to blood loss and hemorrhage in the adult organism. Erythropoietin is synthesized by the kidney and stimulates the proliferation and maturation of bone marrow erythroid precursor cells. The protein is found in the plasma and regulates red cell production by promoting

Product Details

erythroid differentiation and initiating hemoglobin synthesis.

Cross-Reactivity (Details): Human.

Purification: 1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target: EPO

Alternative Name: EPO ([EPO Products](#))

Target Type: Hormone

Background: EP, EPO alpha, EPO, Epoetin, Erythropoietin, MVCD2,Erythropoietin (EPO) (Marker of Placentation Disorders)
Cellular localisation: Cytoplasmic and Cell Surface

Molecular Weight: 37kDa

Gene ID: 2056, 2303

UniProt: [P01588](#)

Pathways: [JAK-STAT Signaling](#), [Hormone Activity](#), [Negative Regulation of intrinsic apoptotic Signaling](#), [Negative Regulation of Transporter Activity](#)

Application Details

Application Notes: Known_Application: ELISA (Use Ab at 2-4 µg/mL for coating) (Order Ab without BSA),
,Immunohistochemistry (Formalin-fixed) (2-4 µg/mL for 30 min at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes), Optimal dilution for a specific application should be determined.
Positive_Control: HepG2 cells. Heart or Kidney.

Restrictions: For Research Use only

Handling

Concentration: 1.0 mg/mL

Buffer: Purified Ab WITHOUT BSA and Azide at 1.0mg/ml

Preservative: Azide free

Handling

Storage:	-20 °C,-80 °C
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.
Expiry Date:	24 months