

Datasheet for ABIN6937161 anti-EBV LMP1 antibody



Overview

Quantity:	100 µg
Target:	EBV LMP1
Reactivity:	Epstein-Barr Virus (EBV)
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This EBV LMP1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (IHC), Staining Methods (StM)

Product Details

Immunogen:	Recombinant fusion protein containing the sequence of bacterial beta-galactosidase and the carboxyl half of EBV-encoded LMP
Clone:	CS4
Isotype:	lgG1 kappa
Specificity:	This antibody is a mixture of four different monoclonal antibodies. This antibody is specific to 60 kDa latent membrane protein (LMP-1) encoded by the BNLF1 gene of the EBV. Each clone reacts with different epitopes on the hydrophilic C-terminus of the cytoplasmic domain of LMP-1. This antibody stains strongly with EBV-positive lymphoblastoid cell lines and EBV infected B cell immunoblasts in infectious mononucleosis. EBV, also designated human herpesvirus 4 (HHV-4), is a member of the herpesvirus family and is one of the most common human viruses. EBV infects B cells and, though often asymptomatic, it can cause infectious mononucleosis, a disease characterized by fatigue, fever, sore throat and muscle soreness.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN6937161 | 07/25/2024 | Copyright antibodies-online. All rights reserved.

Product Details

Purification:

Purified by Protein A/G

Target Details

i di got 2 otalio	
Target:	EBV LMP1
Alternative Name:	Epstein-Barr Virus (LMP-1) (EBV LMP1 Products)
Target Type:	Viral Protein
Molecular Weight:	60kDa
Application Details	
Application Notes:	Positive Control: EBV-infected cells. Tissues.
	Known Application: Immunofluorescence (0.5-1.0 µg/mL), Western Blot (0.5-1.0 µg/mL),
	Immunohistochemistry (Formalin-fixed) (0.5-1.0 μ g/mL for 30 min at RT)(Staining of formalin-
	fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min
	followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be
	determined.
Restrictions:	For Research Use only
Handling	
Buffer:	10 mM PBS without BSA and without Azide.
Preservative:	Azide free
Storage:	4 °C,-80 °C
Storage Comment:	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody
	is stable for 24 months.
Expiry Date:	24 months