## antibodies - online.com







## anti-EPX antibody



( )	11/	IN	/ie	A .
	/ // <del> </del>	۱ ات	/   (−	' \/\/

Quantity:	100 μg	
Target:	EPX	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This EPX antibody is un-conjugated	
Application:	ELISA, Coating (Coat)	

Human eosinophils from a patient with hypereosinophilic syndrome

## **Product Details**

Immunogen:

Clone:	AHE-1
Isotype:	IgG1 kappa
Specificity:	Peripheral blood granulocytes are classified into neutrophils, basophils and eosinophils according to the staining characteristics of their cytoplasmic granules. Granule proteins are released by physiologic and pharmacologic stimuli and play important roles in both normal and pathological host immune responses. Eosinophil major basic protein and eosinophil peroxidase (EPX) are granule proteins specific to the eosinophil. AHE-1 recognizes human EPX, a granule protein specific to eosinophils. It does not cross-react with eosinophil major basic protein, elastase, cathepsin G, esterase N, thrombin, plasmin, kallikrein, lactoferrin, or transferrin. This MAb stains eosinophils only and does not stain other peripheral blood cells, including platelets, neutrophils, monocytes, lymphocytes or red blood cells. Human EPX gene product can form a
	tetramer of two light chains and two heavy chains. Other peroxidase family members include

## **Product Details**

- Todaot Botano		
	myeloperoxidase (MPO), lactoperoxidase (LPO), and thyroid peroxidase (TPO).	
Cross-Reactivity (Details):	Human.	
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.	
Target Details		
Target:	EPX	
Alternative Name:	EPX (EPX Products)	
Background:	Eosinophil peroxidase heavy chain, EPER, EPO, EPP, EPX, EPX PEN, PERE, Eosinophil Peroxidase (EPX)	
	Cellular localisation: Cytoplasmic, granular	
Molecular Weight:	70kDa (55kDa-Heavy Chain, 15kDa-Light Chain)	
Gene ID:	8288, 279259	
UniProt:	P11678	
Application Details		
Application Notes:	Known_Application: ELISA (For coating, order antibody without BSA),Optimal dilution for a	
	specific application should be determined.	
	Positive_Control: MCF-7 cells. Tonsil.	
Restrictions:	For Research Use only	
Handling		
Concentration:	1.0 mg/mL	
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.	
Preservative:	Azide free	
Storage:	-20 °C,-80 °C	
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.	
Expiry Date:	24 months	