

Datasheet for ABIN6937360
anti-GZMB antibody (AA 73-187)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	GZMB
Binding Specificity:	AA 73-187
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This GZMB antibody is un-conjugated
Application:	Immunohistochemistry (IHC), ELISA, Coating (Coat)

Product Details

Immunogen:	Recombinant fragment of human GZMB protein (around aa 73-187) (exact sequence is proprietary)
Clone:	GZMB-2403
Isotype:	IgG2b kappa
Specificity:	Granzyme B is a member of the granule serine protease family stored specifically in NK cells or cytotoxic T cells. Cytolytic T lymphocytes (CTL) and natural killer (NK) cells share the ability to recognize, bind, and lyse specific target cells. They are thought to protect their host by lysing cells bearing on their surface 'nonself' antigens, usually peptides or proteins resulting from infection by intracellular pathogens. Granzyme B is crucial for the rapid induction of target cell apoptosis by CTLs in the cell-mediated immune response. Granzyme B is useful as a marker in the identification of NK/T-cell lymphomas. High percentages of cytotoxic T-cells have been

Product Details

shown to be an unfavorable prognostic indicator in Hodgkin's Disease.

Cross-Reactivity (Details): Human.

Purification: 1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target: GZMB

Alternative Name: GZMB ([GZMB Products](#))

Background: Cathepsin G-like 1, CCPI, CGL1, CSPB, CTLA-1, CTSG1, Cytotoxic serine protease B, Cytotoxic T lymphocyte associated serine esterase 1, Cytotoxic T-lymphocyte proteinase 2, Fragmentin-2, Human lymphocyte protein (Hlp), Lymphocyte protease, SECT, T-cell serine protease 1-3E, Granzyme B (NK/T-Cell Lymphoma Marker)
Cellular localisation: Cytoplasmic

Molecular Weight: 29-32kDa

Gene ID: 3002, 1051

UniProt: [P10144](#)

Pathways: [Apoptosis](#), [Caspase Cascade in Apoptosis](#)

Application Details

Application Notes: Known_Application: ELISA (For coating use Ab at 1-5ug/ml, order Ab without BSA) Optimal dilution for a specific application should be determined.
Positive_Control: A431 cells. Human tonsil, spleen or Hodgkin's Lymphoma tissue (IHC).

Restrictions: For Research Use only

Handling

Concentration: 1.0 mg/mL

Buffer: Prepared in 10 mM PBS, WITHOUT BSA and Azide.

Preservative: Azide free

Storage: -20 °C, -80 °C

Storage Comment: Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.

Expiry Date: 24 months