

Datasheet for ABIN6937365
anti-TIM3 antibody (AA 22-202)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	TIM3 (TIM 3)
Binding Specificity:	AA 22-202
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This TIM3 antibody is un-conjugated
Application:	Immunohistochemistry (Formalin-fixed Sections) (IHC (f))

Product Details

Immunogen:	Recombinant fragment of human TIM3 protein (around aa 22-202) (exact sequence is proprietary)
Clone:	HAVCR2-192
Isotype:	IgG2a kappa
Specificity:	TIMs are type I transmembrane glycoproteins with one Ig-like V-type domain and a Ser/Thr-rich mucin stalk. TIM-3 is expressed on the surface of effector T cell tonsil, lymph node or spleen. Is (CD4+Th1 and CD8+Tc1) but not on helper T cells (CD4+Th2 and CD8+Tc2). In chronic inflammation, autoimmune disorders, and some cancers, TIM-3 is upregulated on several other hematopoietic cell types. The Ig domain of TIM-3 interacts with a ligand on resting but not activated Th1 and Th2 cells. The glycosylated Ig domain of TIM-3 binds cell-associated galectin-9. This induces TIM-3 Tyr phosphorylation and pro-apoptotic signaling. TIM-3

Product Details

functions as a negative regulator of Th1 cell activity. Its blockade results in increased IFN-
gamma production, Th1 cell proliferation and cytotoxicity, regulatory T cell development, and
increases in macrophage and neutrophil infiltration into sites of inflammation.

Cross-Reactivity (Details):	Human.
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target:	TIM3 (TIM 3)
Alternative Name:	HAVCR2 (TIM 3 Products)
Target Type:	Virus
Background:	CD366, HAVR2, Hepatitis A virus cellular receptor 2 (HAVCR2), Kidney injury molecule 3 (KIM3), T-cell immunoglobulin and mucin domain-containing protein 3, T-cell immunoglobulin mucin receptor 3, T-cell membrane protein 3, TIM3, TIMD3,TIM3 / HAVCR2 / CD366 (Effector T-Cell Marker) Cellular localisation: Cell Surface
Molecular Weight:	60kDa
Gene ID:	84868, 710500
UniProt:	Q8TDQ0
Pathways:	Regulation of Lipid Metabolism by PPARalpha , Cancer Immune Checkpoints

Application Details

Application Notes:	Known_Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined. Positive_Control: HepG2 or Daudi cells. Human Tonsil, lymph node or spleen.
Restrictions:	For Research Use only

Handling

Concentration:	1.0 mg/mL
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.

Handling

Preservative:	Azide free
Storage:	-20 °C,-80 °C
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.
Expiry Date:	24 months