

Datasheet for ABIN6937593
anti-CD11c antibody (AA 637-827)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	CD11c (ITGAX)
Binding Specificity:	AA 637-827
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD11c antibody is un-conjugated
Application:	ELISA, Coating (Coat)

Product Details

Immunogen:	Recombinant fragment (aa 637-827) of human ITGAX protein (exact sequence is proprietary)
Clone:	ITGAX-1243
Isotype:	IgG2b kappa
Specificity:	Recognizes a protein of 145 kDa, identified as CD11c. CD11c (ITGAX), a member of the leukointegrin family, shares the same beta subunit with other members of the leukocyte adhesion molecule family, which includes CD11a (LFA-1), CD11b (MAC-1) and CD11d (ITGAD), but has a unique alpha chain. CD11c has been shown to play a role in phagocytosis, cell migration, and cytokine production by monocytes/macrophages as well as induction of T-cell proliferation by Langerhans cells. CD11c is expressed prominently on the plasma membranes of monocytes, tissue macrophages, NK cells, and most dendritic cells (DCs). A lower level of expression is also observed on neutrophils as a result of its high level of expression on most

Product Details

	DCs. An antibody to CD11c may aid in identification of lesions with histiocytic origin. It may also been used as a marker for hairy cell leukemia in paraffin-embedded tissues.
Cross-Reactivity (Details):	Human.
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target:	CD11c (ITGAX)
Alternative Name:	ITGAX (ITGAX Products)
Background:	CD11 antigen-like family member C, Complement component 3 receptor 4 subunit, Integrin alpha-X, integrin, alpha X (antigen CD11C (p150), alpha polypeptide), Leu M5 alpha subunit, Leukocyte adhesion glycoprotein p150 95 alpha chain, Myeloid membrane antigen alpha subunit, p150/95,CD11c (Dendritic Cell Marker) Cellular localisation: Cell Surface
Molecular Weight:	145kDa
Gene ID:	3687, 248472
UniProt:	P20702
Pathways:	Complement System , Activated T Cell Proliferation , Integrin Complex

Application Details

Application Notes:	Known_Application: ELISA (Use Ab at 2-4 µg/mL for coating) (Order Ab without BSA), Optimal dilution for a specific application should be determined. Positive_Control: THP-1 cells.
Restrictions:	For Research Use only

Handling

Concentration:	1.0 mg/mL
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.
Preservative:	Azide free
Storage:	-20 °C,-80 °C
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.

Handling

Expiry Date: 24 months