

Datasheet for ABIN6937688

Recombinant anti-Cytokeratin 18 antibody[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	Cytokeratin 18 (KRT18)
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This Cytokeratin 18 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Flow Cytometry (FACS), Immunohistochemistry (Formalin-fixed Sections) (IHC (f)), Immunostaining (ISt)

Product Details

Immunogen:	Recombinant human full-length KRT18 protein
Clone:	KRT18-2808R
Isotype:	IgG
Specificity:	This MAb reacts with a wide variety of simple epithelia. It does not react with stratified squamous epithelia. It reacts with epithelial tumors of the gastrointestinal tract, lung, breast, pancreas, ovary, and thyroid. Cytokeratin 18, which belongs to the type A (acidic) subfamily of low molecular weight keratins, exists in combination with cytokeratin 8. It is reported that tissues from gastrointestinal tract are positive for both cytokeratin 8 and 18 but do not contain cytokeratin 14. Tissues from gastrointestinal tract, respiratory tract and urogenital tract, as well as endocrine and exocrine tissues and mesothelial cells are positive for cytokeratin 18.

Product Details

Cross-Reactivity (Details): Human. Does not react with Dog.

Purification: 1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target: Cytokeratin 18 (KRT18)

Alternative Name: KRT18 ([KRT18 Products](#))

Background: Cell Proliferation-inducing Gene 46 Protein, CK18, CYK18Cytokeratin Endo B, K18, Keratin-18, Kerd, KRT18,Cytokeratin 18 (KRT18)
Cellular localisation: Cytoplasmic

Molecular Weight: 45kDa

Gene ID: 3875, 406013

UniProt: [P05783](#)

Pathways: [Apoptosis](#), [Caspase Cascade in Apoptosis](#)

Application Details

Application Notes: Known_Application: Flow Cytometry (1-2 µg/million cells), Immunofluorescence (1-2 µg/mL),Western Blot (1-2 µg/mL),Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 min at RT)(Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0 for 45 min at 95°C followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.
Positive_Control: HCT116, MCF-7, HeLa, A431 cells. Breast Cancer.

Restrictions: For Research Use only

Handling

Concentration: 1.0 mg/mL

Buffer: Prepared in 10 mM PBS, WITHOUT BSA and Azide.

Preservative: Azide free

Storage: -20 °C,-80 °C

Storage Comment: Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.

Expiry Date: 24 months