

Datasheet for ABIN6937788

Recombinant anti-LMO2 antibody (AA 23-140)[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	LMO2
Binding Specificity:	AA 23-140
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This LMO2 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunofluorescence (IF)

Product Details

Immunogen:	Recombinant fragment (around aa 23-140) human LMO2 protein (exact sequence is proprietary)
Clone:	LMO2-3147R
Isotype:	IgG
Specificity:	The LMO2 protein has a central and crucial role in hematopoietic development and is highly conserved. It has a particular function in normal and lymphatic endothelial cells involving the regulation of angiogenesis and lymph-angiogenesis. Immunohistochemical studies have also demonstrated expression of LMO2 in both normal germinal center B-cells and germinal center-derived B-cell lymphomas, including follicular lymphoma and diffuse large B-cell lymphoma. The use of anti-LMO2 is valuable as a tool in the identification of lymphomas of B-cell origin.

Product Details

LMO2 is useful in differentiating follicular lymphoma (LMO2+) from nodal marginal zone lymphoma (LMO2-). It also is positive in Hodgkin's and Burkitt's lymphomas.

Cross-Reactivity (Details): Human.

Purification: 1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target: LMO2

Alternative Name: LMO2 ([LMO2 Products](#))

Background: Cysteine-rich protein TTG-2, LIM domain only protein 2, LMO-2, RBTN L1, RBTN2, Rhombotin like 1, Rhombotin-2 (RHOM2), T-cell translocation protein 2, TTG2, LMO2 (B-Cell Marker)
Cellular localisation: Nuclear

Molecular Weight: 24kDa

Gene ID: 4005, 34560

UniProt: [P25791](#)

Pathways: [Chromatin Binding](#)

Application Details

Application Notes: Known_Application: Flow Cytometry (1-2ug/million cells); ,Immunofluorescence (1-2ug/ml);,Optimal dilution for a specific application should be determined.
Positive_Control: K562, Ramos or Raji cells. Placenta, Pancreas or Hodgkin's lymphoma

Restrictions: For Research Use only

Handling

Concentration: 1.0 mg/mL

Buffer: Prepared in 10 mM PBS, WITHOUT BSA and Azide.

Preservative: Azide free

Storage: -20 °C,-80 °C

Storage Comment: Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.

Expiry Date: 24 months