

Datasheet for ABIN6938024

Recombinant anti-CD56 antibody[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	CD56 (NCAM1)
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This CD56 antibody is un-conjugated
Application:	Flow Cytometry (FACS), Immunohistochemistry (Formalin-fixed Sections) (IHC (f))

Product Details

Immunogen:	Recombinant full-length human NCAM1 (CD56) protein
Clone:	NCAM1-2217R
Isotype:	IgG
Specificity:	This MAb reacts with an extracellular domain (close to transmembrane) of CD56/NCAM. Three isoforms of neural cell adhesion molecule (NCAM) are produced by differential splicing of the RNA transcript from a single gene. The 135 kDa isoform is the basic molecule, which is glycosylated or sialylated to produce the mature species. Anti-CD56 recognizes two proteins of the neural cell adhesion molecule, the basic molecule expressed on most neuroectodermally derived tissues and neoplasms (e.g. retinoblastoma, medulloblastomas, astrocytomas, neuroblastomas, and small cell carcinomas). It is also expressed on some mesodermally derived tumors (rhabdomyosarcoma). Anti-CD56 plays an important role in the diagnosis of

Product Details

	nodal and nasal NK/T-cell lymphomas.
Cross-Reactivity (Details):	Human.
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target:	CD56 (NCAM1)
Alternative Name:	NCAM1 (NCAM1 Products)
Background:	NCAM, Leu-19, NKH1, MSK39, NCAM120, NCAM140, NCAM180, Neural Cell Adhesion Molecule,CD56 / NCAM1 / NKH1 (Neuronal Cell Marker) Cellular localisation: Cell Surface and Cytoplasmic
Molecular Weight:	180, 145 and 125kDa
Gene ID:	4684, 503878
UniProt:	P13591 , P13592

Application Details

Application Notes:	Known_Application: Flow Cytometry (0.5-1 µg/million cells), Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes), Optimal dilution for a specific application should be determined. Positive_Control: PBMCs, Colon or Small Intestine.
Restrictions:	For Research Use only

Handling

Concentration:	1.0 mg/mL
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.
Preservative:	Azide free
Storage:	-20 °C,-80 °C
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.
Expiry Date:	24 months