

Datasheet for ABIN6938070 anti-NKX2-8 antibody (AA 10-123)



Overview

Overview	
Quantity:	100 μg
Target:	NKX2-8
Binding Specificity:	AA 10-123
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This NKX2-8 antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (IHC), Coating (Coat), Flow Cytometry (FACS)
Product Details	
Immunogen:	Human recombinant NKX2.8 protein fragment (around aa 10-123) (exact sequence is proprietary)
Clone:	NKX28-2548
Isotype:	IgG1 kappa
Specificity:	The protein encoded by this gene is a homeobox-containing developmental regulator associated with liver development. The encoded protein binds to the alpha-fetoprotein (AFP) gene promoter and increases the expression of AFP. This gene is overexpressed in some lung cancers and is linked to poor patient survival, possibly due to its resistance to cisplatin. This gene is aberrantly methylated in pancreatic cancer, deleted in squamous cell lung carcinomas, and acts as a tumor suppressor in esophageal cancer. Mutations in this gene may also be a

Product Details

Troduct Details	
	cause of neural tube defects.
Cross-Reactivity (Details):	Human.
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.
Target Details	
Target:	NKX2-8
Alternative Name:	NKX2-8 (NKX2-8 Products)
Background:	Homeobox protein NK 2 homolog H, Homeobox protein NK-2 homolog H, Homeobox protein
	Nkx 2.8, Homeobox protein Nkx-2.8, NK 2 homolog 8, NK 2 homolog H, NK2 homeobox 8, NK2
	transcription factor related, locus 8, Nkx2 9, Nkx2-8, NKX2.8, NKX28_HUMAN, NKX2G, NKX2H,
	OTTHUMP00000027959,NKX2.8
	Cellular localisation: Nuclear and Cytoplasm.
Molecular Weight:	26kDa
Gene ID:	26257, 234763
UniProt:	015522
Application Details	
Application Notes:	Known_Application: Western Blot (2-4 µg/mL), ELISA (For coating, order antibody without BSA),
	Flow Cytometry (1-2 µg/million cells), Western Blot (1-2 µg/mL), Optimal dilution for a specific
	application should be determined.
	Positive_Control: Human Liver, Kidney or Thyroid tissue (IHC).
Restrictions:	For Research Use only
Handling	
Concentration:	1.0 mg/mL
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.
Preservative:	Azide free
Storage:	-20 °C,-80 °C
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.
Expiry Date:	24 months