

Datasheet for ABIN6938116
anti-PAX2A antibody (AA 223-354)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	PAX2A
Binding Specificity:	AA 223-354
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This PAX2A antibody is un-conjugated
Application:	ELISA, Coating (Coat), Immunohistochemistry (Formalin-fixed Sections) (IHC (f))

Product Details

Immunogen:	Recombinant fragment (around aa 223-354) human PAX2 protein (exact sequence is proprietary)
Clone:	PAX2-1104
Isotype:	IgG2b kappa
Specificity:	Recognizes a protein of 42kDa, which is identified as PAX2. It is a member of the paired box family of transcription factors, which is required for development and proliferation of the kidney, brain, and mullerian organs. PAX2 genes contain a highly conserved DNA sequence within the paired box region, which encodes a DNA-binding domain, enabling PAX proteins to bind the promoters of specific genes to transcriptionally regulate their expression. PAX2 is specifically expressed in the developing central nervous system, eye, ear, and urogenital tract, and is essential for the development of these organs. In normal adult tissues PAX2 was mainly

Product Details

detected in the urogenital system, including kidney, ureteric epithelium, fallopian tube epithelium, ovary and uterus. In tumors, PAX2 has been detected in renal cell carcinomas, Wilms' tumors, nephrogenic adenomas and papillary serous carcinoma of the ovary. PAX2 has been used as a marker for the identification of renal cell carcinoma and ovarian carcinoma by immunohistochemistry.

Cross-Reactivity (Details):	Human.
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target:	PAX2A
Alternative Name:	PAX2 (PAX2A Products)
Background:	Paired box gene 2, paired box homeotic gene 2, paired box protein 2, Paired box protein Pax-2, Paired box protein Pax2,PAX2 (Renal Cell & Ovarian Carcinoma Marker) Cellular localisation: Nuclear
Molecular Weight:	42kDa
Gene ID:	5076, 155644
UniProt:	Q02962
Pathways:	Carbohydrate Homeostasis , Stem Cell Maintenance , Tube Formation

Application Details

Application Notes:	Known_Application: ELISA (For coating, order Ab without BSA),Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes), Optimal dilution for a specific application should be determined. Positive_Control: NAMALWA cells. Fetal kidney or Renal Cell Carcinoma (RCC) or Ovarian Carcinoma.
Restrictions:	For Research Use only

Handling

Concentration:	1.0 mg/mL
----------------	-----------

Handling

Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.
Preservative:	Azide free
Storage:	-20 °C,-80 °C
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.
Expiry Date:	24 months