

Datasheet for ABIN6938213

Recombinant anti-POMC antibody (AA 25-39)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	POMC
Binding Specificity:	AA 25-39
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This POMC antibody is un-conjugated
Application:	ELISA, Coating (Coat)

Product Details

Immunogen:	Synthetic peptide corresponding to aa25-39 of human ACTH (NGAEDESAEAFPLEF)
Clone:	CLIP-2859R
Isotype:	IgG
Specificity:	ACTH (same as Corticotropin) is a 39 amino acid active peptide produced by the anterior pituitary. This MAb is specific to CLIP (aa25-39 of ACTH); does not react with Synacthen (aa1-24 of ACTH). POMC (pro-opiomelanocortin or corticotropin-lipotropin) is a 267 amino acid polypeptide hormone precursor that goes through extensive, tissue-specific posttranslational processing by convertases. POMC is cleaved into ten hormone chains named NPP, ACTH, alpha-MSH (Melanocyte Stimulating Hormone), beta-MSH, gamma-MSH, CLIP (corticotropin-like intermediary peptide), Lipotropin-beta, Lipotropin-gamma, beta-endorphin and Met-

Product Details

enkephalin. ACTH is also produced by cells of immune system (T-cells, B-cells, and macrophages) in response to stimuli associated with stress. Anti-ACTH is a useful marker in classification of pituitary tumors and the study of pituitary disease. It reacts with ACTH-producing cells (corticotrophs). It also may react with other tumors (e.g. some small cell carcinomas of the lung) causing paraneoplastic syndromes by secreting ACTH. '

Cross-Reactivity (Details):	Human.
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target:	POMC
Alternative Name:	POMC (POMC Products)
Background:	Adrenocorticotropin, alpha or beta or gamma Melanocyte Stimulating Hormone (MSH) or Melanotropin, beta-Endorphin, beta or gamma Lipotropin (LPH), CLIP, Met Enkephalin, POC, POMC, ACTH (Adrenocorticotrophic Hormone) (C-Terminus) Cellular localisation: Cytoplasmic
Molecular Weight:	ACTH is ~5kDa, and the POMC precursor is ~30kDa. The molecular weight of POMC depends upon isoform variation and post-translational modifications.
Gene ID:	5443, 1897
UniProt:	P01189
Pathways:	Metabolism of Steroid Hormones and Vitamin D , Peptide Hormone Metabolism , Hormone Activity , Feeding Behaviour

Application Details

Application Notes:	Known_Application: ELISA (For coating, order Ab without BSA), Optimal dilution for a specific application should be determined. Positive_Control: Normal pituitary gland or pituitary tumor.
Restrictions:	For Research Use only

Handling

Concentration:	1.0 mg/mL
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.

Handling

Preservative:	Azide free
Storage:	-20 °C,-80 °C
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.
Expiry Date:	24 months