antibodies -online.com





Datasheet for ABIN6938354

anti-SAA antibody



	Θ_{W}

Quantity:	100 μg
Target:	SAA
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This SAA antibody is un-conjugated
Application:	Immunostaining (ISt), Immunohistochemistry (Formalin-fixed Sections) (IHC (f))
Product Details	
Immunogen:	Recombinant full-length human serum amyloid A (SAA) protein
Clone:	SAA-326
Isotype:	IgG2a kappa
Specificity:	This antibody reacts with natural and recombinant human Serum Amyloid A (SAA) and does
	not cross-react with other human cytokines or growth factors. Human SAA proteins are a group
	of apo-lipoproteins found predominantly in the high-density lipoprotein (HDL) fraction of
	plasma. SAA is a major acute-phase protein and precursor to amyloid A protein, which is the
	major constituent of the fibril deposits of reactive amyloidosis. SAA is secreted in large
	amounts by the liver during microbial infections or inflammatory diseases.
Cross-Reactivity (Details):	Human.
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target:	SAA
Alternative Name:	SAA (SAA Products)
Background:	Amyloid fibril protein AA, Amyloid fibrils, PIG4, Serum amyloid A protein (SAA), Serum amyloid A1 isoform 1 (SAA1), Serum amyloid A1 isoform 2 (SAA2), Tumor protein p53 inducible protein 4 (TP53I4), Serum Amyloid A Cellular localisation: Cytoplasmic & extracellular (secreted)
Molecular Weight:	12kDa
Gene ID:	6288, 6289, 632144
UniProt:	P02735, P0DJI8

Application Details

Application Notes:	Known_Application: Immunohistochemistry (Formalin-fixed) (1-2 μg/mL for 30 minutes at
	RT),(Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10 mM Tris
	with 1 mM EDTA, pH 9.0 for 45 min at 95°C followed by cooling at RT for
	20 minutes),Optimal dilution for a specific application should be determined.
	Positive_Control: Caki-1 cells. Pancreas or Liver.
Restrictions:	For Research Use only

Handling

Concentration:	1.0 mg/mL	
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.	
Preservative:	Azide free	
Storage:	-20 °C,-80 °C	
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.	
Expiry Date:	24 months	