

Datasheet for ABIN6938384 **anti-Melanoma gp100 antibody**



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	Melanoma gp100 (PMEL)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Melanoma gp100 antibody is un-conjugated
Application:	Immunohistochemistry (Formalin-fixed Sections) (IHC (f))

Product Details

Immunogen:	Membranes from a human melanoma metastasis
Clone:	NKI-beteb
Isotype:	IgG2b kappa
Specificity:	By immunohistochemistry, it specifically recognizes a protein in melanocytes and melanomas. This MAb reacts with junctional and blue nevus cells and variably with fetal and neonatal melanocytes. Intradermal nevi, normal adult melanocytes, and non-melanocytic cells are negative. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin. This Mab labels formalin-fixed, paraffin-embedded melanomas and other tumors showing melanocytic differentiation.
No Cross-Reactivity:	Rat (Rattus)
Cross-Reactivity (Details):	Human. Horse. Does not react with Rat.
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target:	Melanoma gp100 (PMEL)
Alternative Name:	SILV (PMEL Products)
Background:	95 kDa melanocyte-specific secreted glycoprotein, M-beta, Melanocyte lineage specific antigen GP100, Melanocyte protein Pmel 17, Melanoma associated ME20 antigen, Melanosomal matrix protein17, p100, p26, PMEL17, Premelanosome protein, Secreted melanoma-associated ME20 antigen, SILV, Silver homolog,gp100 / Melanosome / PMEL17 / SILV (Melanoma Marker) Cellular localisation: Cytoplasmic
Molecular Weight:	90-100kDa
Gene ID:	6490, 95972
UniProt:	P40967

Application Details

Application Notes:	Known_Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes), Optimal dilution for a specific application should be determined. Positive_Control: SK-MEL-28 cells. Melanoma.
Restrictions:	For Research Use only

Handling

Concentration:	1.0 mg/mL
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.
Preservative:	Azide free
Storage:	-20 °C,-80 °C
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.
Expiry Date:	24 months