

Datasheet for ABIN6938393

Recombinant anti-Melanoma gp100 antibody



Overview

Quantity:	100 μg
Target:	Melanoma gp100 (PMEL)
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This Melanoma gp100 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Formalin-fixed Sections) (IHC (f))
Product Details	
Immunogen:	Recombinant full-length human gp100 protein
Immunogen: Clone:	Recombinant full-length human gp100 protein PMEL-1825R
Clone:	PMEL-1825R
Clone:	PMEL-1825R
Clone:	PMEL-1825R IgG The gp100 Molecule is a 100 kDa glycosylated protein that is cleaved into a small (26 kDa)
Clone:	PMEL-1825R IgG The gp100 Molecule is a 100 kDa glycosylated protein that is cleaved into a small (26 kDa) carboxy-terminal fragment and a larger amino- terminal section (60 64 kDa), which is
Clone:	PMEL-1825R IgG The gp100 Molecule is a 100 kDa glycosylated protein that is cleaved into a small (26 kDa) carboxy-terminal fragment and a larger amino- terminal section (60 64 kDa), which is subsequently cleaved to generate 26 kDa and 34 38 kDa fragments. By immunohistochemistry,
Clone:	PMEL-1825R IgG The gp100 Molecule is a 100 kDa glycosylated protein that is cleaved into a small (26 kDa) carboxy-terminal fragment and a larger amino- terminal section (60 64 kDa), which is subsequently cleaved to generate 26 kDa and 34 38 kDa fragments. By immunohistochemistry, it specifically recognizes a protein in melanocytes and melanomas. This MAb reacts with
Clone:	PMEL-1825R IgG The gp100 Molecule is a 100 kDa glycosylated protein that is cleaved into a small (26 kDa) carboxy-terminal fragment and a larger amino- terminal section (60 64 kDa), which is subsequently cleaved to generate 26 kDa and 34 38 kDa fragments. By immunohistochemistry, it specifically recognizes a protein in melanocytes and melanomas. This MAb reacts with junctional and blue nevus cells and variably with fetal and neonatal melanocytes. Intradermal
Clone:	IgG The gp100 Molecule is a 100 kDa glycosylated protein that is cleaved into a small (26 kDa) carboxy-terminal fragment and a larger amino- terminal section (60 64 kDa), which is subsequently cleaved to generate 26 kDa and 34 38 kDa fragments. By immunohistochemistry, it specifically recognizes a protein in melanocytes and melanomas. This MAb reacts with junctional and blue nevus cells and variably with fetal and neonatal melanocytes. Intradermal nevi, normal adult melanocytes, and non-melanocytic cells are negative. It does not stain tumor

Product Details

	sarcomas using H & E stains alone. It is also difficult to differentiate melanoma from spindle cell carcinomas and various types of mesenchymal neoplasms. This MAb stains fetal and neonatal melanocytes, junctional and blue nevus cells, and malignant melanoma. This MAb also stains Angiomyolipoma (PEComa).
No Cross-Reactivity:	Dog (Canine), Rat (Rattus)
Cross-Reactivity (Details):	Human. Does not react with dog and rat
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.
Target Details	
Target:	Melanoma gp100 (PMEL)
Alternative Name:	SILV (PMEL Products)
Background:	95 kDa melanocyte-specific secreted glycoprotein, M-beta, Melanocyte lineage specific antiger GP100, Melanocyte protein Pmel 17, Melanosomal matrix protein17, p100, p26, SILV, Silver homolog,gp100 / Melanosome / PMEL17 / SILV (Melanoma Marker) Cellular localisation: Cytoplasm
Molecular Weight:	26-100kDa
Gene ID:	6490, 95972
UniProt:	P40967
Application Details	
Application Notes:	Known_Application: Western Blot (1-2 μ g/mL), ,Immunohistochemistry (Formalin-fixed) (1-2 μ g/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined. Positive_Control: COLO-38 or SK-MEL-28 cells. Human melanoma.
Restrictions:	For Research Use only
Handling	
Concentration:	1.0 mg/mL
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.
Preservative:	Azide free

Handling

Storage:	-20 °C,-80 °C
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.
Expiry Date:	24 months