

Datasheet for ABIN6938397

Recombinant anti-GLUT1 antibody (AA 203-305)[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	GLUT1 (SLC2A1)
Binding Specificity:	AA 203-305
Reactivity:	Human
Host:	Mouse
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This GLUT1 antibody is un-conjugated
Application:	ELISA, Coating (Coat), Immunohistochemistry (Formalin-fixed Sections) (IHC (f))

Product Details

Immunogen:	Recombinant fragment of human GLUT1 protein (around aa 203-305) (exact sequence is proprietary)
Clone:	RGLUT1-2476
Isotype:	IgG1 kappa
Specificity:	Recognizes a protein of 55 kDa, which is identified as GLUT-1. Glucose transporters are integral membrane glycoproteins involved in transporting glucose into most cells. There are many types of glucose transport carrier proteins, designated as Glut-1 to Glut-12. Glut-1 is a major glucose transporter in the mammalian blood-brain barrier. It is expressed in high density on the membranes of human erythrocytes and the brain capillaries that comprise the blood-brain barrier. Glut-1 is expressed at variable levels in many human tissues. Overexpression of Glut-1

Product Details

has been linked to tumor progression or poor survival of patients with carcinomas of the colon, breast, cervical, lung, bladder and mesothelioma. Glut-1 is a sensitive and specific marker for the differentiation of malignant mesothelioma (positive) from reactive mesothelium (negative).

Cross-Reactivity (Details): Human.

Purification: 1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target: GLUT1 (SLC2A1)

Alternative Name: SLC2A1 ([SLC2A1 Products](#))

Background: Erythrocyte/hepatoma glucose transporter, Glucose transporter type-1, GLUT1, GLUT1DS, GLUTB, GT1, GTG1, Gtg3, HepG2 glucose transporter, PED, RATGTG1, Solute carrier family 2, Solute carrier family 2, facilitated glucose transporter member 1 (SLC2A1), GLUT-1 (Tumor Progression and Mesothelioma Marker)
Cellular localisation: Cell Surface

Molecular Weight: 55kDa

Gene ID: 6513, 473721

UniProt: [P11166](#)

Pathways: [Sensory Perception of Sound](#), [Dicarboxylic Acid Transport](#), [Warburg Effect](#)

Application Details

Application Notes: Known_Application: ELISA (For coating use Ab at 1-2 µg/mL order Ab without BSA), Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes), Optimal dilution for a specific application should be determined.
Positive_Control: K562, A431, MDA-MB-231 cells or erythrocytes. Mesothelioma or breast, colon and ovarian carcinoma.

Restrictions: For Research Use only

Handling

Concentration: 1.0 mg/mL

Handling

Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.
Preservative:	Azide free
Storage:	-20 °C,-80 °C
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.
Expiry Date:	24 months