antibodies -online.com





Recombinant anti-SPTA1 antibody (AA 356-475)



()	ve	K\ /		A .
	\cup	1 V/	Щ.	V۷

Cross-Reactivity (Details):

Human.

Quantity:	100 μg	
Target:	SPTA1	
Binding Specificity:	AA 356-475	
Reactivity:	Human	
Host:	Mouse	
Antibody Type:	Recombinant Antibody	
Clonality:	Monoclonal	
Conjugate:	This SPTA1 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (Formalin-fixed Sections) (IHC (f))	
Product Details		
Immunogen:	Recombinant fragment of human SPTA (around aa356-475) (exact sequence is proprietary)	
Clone:	RSPTA1-1832	
Isotype:	IgG1 kappa	
Specificity:	Spectrin is an actin crosslinking and molecular scaffold protein that links the plasma membrane to the actin cytoskeleton, and functions in the determination of cell shape, arrangement of transmembrane proteins, and organization of organelles. Spectrins funct membrane organizers and stabilizers, composed of non-homologous) are present in oth somatic cells. The spectrin tetramers in erythrocytes act as barriers to lateral diffusion, by spectrin dimers seem to lack this function.	

Product Details Purification: 1.0mg/ml of Ab purified from Bioreactor by Protein A/G. **Target Details** SPTA1 Target: SPTA1 (SPTA1 Products) Alternative Name Background: Alpha I spectrin, Elliptocytosis 2 (EL2), Erythrocyte alpha spectrin, Erythroid alpha-spectrin, HPP, HS3, Spectrin alpha erythrocytic 1, SPH3, SPTA1, Spectrin Alpha 1 (Erythrocyte Marker) Cellular localisation: Cytoplasmic and Cell Surface 240kDa Molecular Weight: Gene ID: 6708, 119825 UniProt: P02549 Pathways: Regulation of Actin Filament Polymerization **Application Details** Known_Application: Western Blot (1-2 μg/mL), Immunohistochemistry (Formalin-fixed) (1-2 μ **Application Notes:** g/mL for 30 minutes at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes), Optimal dilution for a specific application should be determined. Positive_Control: K562. HeLa, Jurkat cells. Placenta or Tonsil. Restrictions: For Research Use only Handling Concentration: 1.0 mg/mL Buffer: Prepared in 10 mM PBS, WITHOUT BSA and Azide. Preservative: Azide free -20 °C,-80 °C Storage: Storage Comment: Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.

24 months

Expiry Date: