

Datasheet for ABIN6938547
anti-Transglutaminase 2 antibody



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	Transglutaminase 2 (TGM2)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Transglutaminase 2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Formalin-fixed Sections) (IHC (f))

Product Details

Immunogen:	Recombinant full-length human TGM2 protein
Clone:	SPM358
Isotype:	IgG2a kappa
Specificity:	Recognizes a 77-85 kDa protein, identified as cellular or tissue transglutaminase II (TGase II). Transglutaminases are enzymes that catalyze the crosslinking of proteins by epsilon-gamma glutamyl lysine isopeptide bonds. While the primary structure of transglutaminases is not conserved, they all have the same amino acid sequence at their active sites and their activity is calcium-dependent. The protein encoded by this gene acts as a monomer, is induced by retinoic acid, and appears to be involved in apoptosis. Finally, the encoded protein is the autoantigen implicated in celiac disease. The identification of transglutaminase as the main antigen of endomysium antibodies allows a new diagnostic approach to celiac disease (CD), a genetic, immunologically mediated small bowel enteropathy that causes malabsorption. TGase II is implicated in programmed cell death, signal transduction, drug-resistance, cell growth,

Product Details

endocytosis, insulin secretion, cell adhesion, cataract formation, and wound healing.

Cross-Reactivity (Details): Human, Monkey, Rabbit, Mouse and Rat.

Purification: 1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target: Transglutaminase 2 (TGM2)

Alternative Name: TGM2 ([TGM2 Products](#))

Background: Protein-glutamine gamma-glutamyltransferase 2, TG(C), TG2, TGase C, TGase H, TGase-2, Tgasell, TGC, TGM2, Tissue type transglutaminase, Transglutaminase 2 C polypeptide, Transglutaminase H, tTG, tTGas, Transglutaminase II (TGM2)
Cellular localisation: Cytoplasm. Cell surface.

Molecular Weight: 77-85kDa

Gene ID: 7052, 517033

UniProt: [P21980](#)

Pathways: [Tube Formation](#), [Thromboxane A2 Receptor Signaling](#)

Application Details

Application Notes: Known_Application: Western Blot (1-2 µg/mL), ,Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95 °C followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined.
Positive_Control: HUVEC cells. Endothelial cells in placenta, liver, brain, or breast carcinoma.
Smooth muscle cells of any origin

Restrictions: For Research Use only

Handling

Concentration: 1.0 mg/mL

Buffer: Prepared in 10 mM PBS, WITHOUT BSA and Azide.

Preservative: Azide free

Storage: -20 °C,-80 °C

Handling

Storage Comment: Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.

Expiry Date: 24 months