## antibodies -online.com





## anti-TOX3 antibody (AA 251-389)



Go to Product page

( )	11/	IN	/ie	A .
	/ // <del> </del>	۱ ات	/   (−	' \/\/

Quantity:	100 μg		
Target:	TOX3		
Binding Specificity:	AA 251-389		
Reactivity:	Human		
Host:	Mouse		
Clonality:	Monoclonal		
Conjugate:	This TOX3 antibody is un-conjugated		
Application:	Western Blotting (WB), ELISA, Coating (Coat)		
Product Details			
Immunogen:	Recombinant fragment of human TOX3 protein (around aa 251-389) (Exact sequence is		
	proprietary)		
Clone:	TOX3-1124		
Isotype:	IgG1 kappa		
Specificity:	It recognizes a 63 kDa protein, which is identified as TOX3. It contains a high mobility group		
	(HMG)-box, which regulates Ca2+-dependent transcription in neurons through interaction with		
	the cAMP-response-element-binding protein (CREB). TOX3 appears to be associated with		
	breast cancer susceptibility and is expressed downstream of a cytoprotective cascade together		
	with CITED1, a transcriptional regulator that does not bind directly to DNA. TOX3 is		
	predominantly expressed in the brain and forms homodimers. TOX3 overexpression protects		
	neuronal cells from cell death caused by endoplasmic reticulum stress or BAX overexpression		

## **Product Details**

	through the induction of anti-apoptotic transcripts and repression of pro-apoptotic transcripts	
Cross-Reactivity (Details):	Human.	
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.	
Target Details		
Target:	TOX3	
Alternative Name:	TOX3 (TOX3 Products)	
Background:	TOX3, CAG trinucleotide repeat-containing gene F9 protein, CAGF9, TOX high mobility group	
	box family member 3, Trinucleotide repeat containing 9 (TNRC9),TOX3 / TNRC9	
	Cellular localisation: Cytoplasmic & Nuclear	
Molecular Weight:	63kDa	
Gene ID:	27324, 460789	
UniProt:	015405	
Pathways:	Chromatin Binding	
Application Details		
Application Notes:	Known_Application: ELISA (For coating, order Ab without BSA), Western Blot (1-2 µg/mL),	
	Optimal dilution for a specific application should be determined.	
	Positive_Control: SH-SY5Y, MDA-MB-435 cells. Brain or breast or gastric Carcinoma.	
Restrictions:	For Research Use only	
Handling		
Concentration:	1.0 mg/mL	
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.	
Preservative:	Azide free	
Storage:	-20 °C,-80 °C	
	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous	
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardou	