

Datasheet for ABIN6938702

anti-Tyrosinase-Related Protein 1 antibody



Overview

Quantity:	100 μg
Target:	Tyrosinase-Related Protein 1 (TYRP1)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Tyrosinase-Related Protein 1 antibody is un-conjugated
Application:	Immunofluorescence (IF), Flow Cytometry (FACS), Western Blotting (WB), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	SK-MEL-23 cells (TA99), Recombinant full-length human TYRP1 protein (TYRP1/807)
Clone:	TA99-TYRP1-807
Isotype:	IgG2a kappa
Specificity:	It reacts with a 75 kDa melanocyte-specific gene product, identified as Tyrosinase-related protein-1 (TRP-1). It is involved in melanin synthesis. TRP1 is present on the melanosomal membranes of melanoma, normal melanocytes and nevi.Recent evidence suggests that TRP-1 is involved in maintaining stability of tyrosinase protein and modulating its catalytic activity. TRP-1 is also involved in maintenance of melanosome ultrastructure and affects melanocyte proliferation and cell death.
Cross-Reactivity (Details):	Human. Mouse.
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details	
Target:	Tyrosinase-Related Protein 1 (TYRP1)
Alternative Name:	TYRP1 (TYRP1 Products)
Background:	5, 6 dihydroxyindole 2 carboxylic acid oxidase, 6-dihydroxyindole-2-carboxylic acid oxidase, Associated with iris pigmentation, CAS2, Catalase B (CATB), DHICA oxidase, Glycoprotein75
	(GP75), Melanoma antigen gp75, Tyrosinase-related protein 1 (TYRP1), TYRRP,Tyrosinase-
	Related Protein-1 (TRP-1) (Melanoma Marker)
	Cellular localisation: Cytoplasmic
Molecular Weight:	75kDa
Gene ID:	7306, 270279
UniProt:	P17643
Application Details	
Application Notes:	Known_Application: Flow Cytometry (1-2 μg/million cells), Immunofluorescence (1-2 μg/mL),
	Western Blot (1-2 μg/mL), Immunohistochemistry (Frozen only) (1-2 μg/mL for 30 minutes at
	RT)Optimal dilution for a specific application should be determined.

Restrictions:	For Research Use only
	Positive_Control: SK-MEL-23, SK-MEL-19, SK-MEL-30, SK-MEL-37 cells. Melanoma.
	RT)Optimal dilution for a specific application should be determined.
	Western Blot (1-2 μ g/mL), Immunohistochemistry (Frozen only) (1-2 μ g/mL for 30 minutes at
Application Notes:	Known_Application: Flow Cytometry (1-2 μg/million cells), Immunofluorescence (1-2 μg/mL),

Handling

Concentration:	1.0 mg/mL
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.
Preservative:	Azide free
Storage:	-20 °C,-80 °C
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.
Expiry Date:	24 months