

Datasheet for ABIN6938703

anti-Tyrosinase-Related Protein 1 antibody (AA 257-377)[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	Tyrosinase-Related Protein 1 (TYRP1)
Binding Specificity:	AA 257-377
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Tyrosinase-Related Protein 1 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), ELISA, Immunohistochemistry (Formalin-fixed Sections) (IHC (f)), Coating (Coat)

Product Details

Immunogen:	A recombinant fragment (around aa 257-377) of human TYRP1 protein (exact sequence is proprietary)
Clone:	TYRP1-3280
Isotype:	IgG2b kappa
Specificity:	It reacts with a 75 kDa melanocyte-specific gene product, identified as Tyrosinase-related protein-1 (TRP-1). It is involved in melanin synthesis. TRP1 is present on the melanosomal membranes of melanoma, normal melanocytes and nevi. Recent evidence suggests that TRP-1 is involved in maintaining stability of tyrosinase protein and modulating its catalytic activity. TRP-1 is also involved in maintenance of melanosome ultrastructure and affects melanocyte proliferation and cell death.

Product Details

Cross-Reactivity (Details):	Human.
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target:	Tyrosinase-Related Protein 1 (TYRP1)
Alternative Name:	TYRP1 (TYRP1 Products)
Background:	5, 6 dihydroxyindole 2 carboxylic acid oxidase, 6-dihydroxyindole-2-carboxylic acid oxidase, Associated with iris pigmentation, CAS2, Catalase B (CATB), DHICA oxidase, Glycoprotein75 (GP75), Melanoma antigen gp75,Tyrosinase-Related Protein-1 (TYRP-1) (Melanoma Marker) Cellular localisation: Cytoplasmic
Molecular Weight:	75kDa
Gene ID:	7306, 270279
UniProt:	P17643

Application Details

Application Notes:	Known_Application: ELISA (Use Ab at 2-4 µg/mL for coating) (Order Ab without BSA), ,Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes), Optimal dilution for a specific application should be determined. Positive_Control: SK-MEL-23, SK-MEL-19, SK-MEL-30, SK-MEL-37 cells. Human skin or melanoma tissues (IHC).
Restrictions:	For Research Use only

Handling

Concentration:	1.0 mg/mL
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.
Preservative:	Azide free
Storage:	-20 °C,-80 °C
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.

Handling

Expiry Date: 24 months