

Datasheet for ABIN6938760
anti-VLDLR antibody (C-Term)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	VLDLR
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This VLDLR antibody is un-conjugated
Application:	Immunostaining (IST), Immunohistochemistry (Formalin-fixed Sections) (IHC (f))

Product Details

Immunogen:	Recombinant human VLDLR fragment from c-terminal
Clone:	VLDLR-1337
Isotype:	IgG1 kappa
Specificity:	VLDLR (very low density lipoprotein receptor) is a member of the LDL receptor gene family, which includes LDL receptor, LRP, megalin, VLDLR and ApoER2. The LDL receptor family is characterized by a cluster of cysteine-rich class A repeats, epidermal growth factor (EGF)-like repeats, YWTD repeats and an O-linked sugar domain. VLDLR associates with RAP (receptor associated protein) during receptor folding, and RAP facilitates the secretion of the extracellular region of VLDLR. VLDLR is thought to mediate the interaction of extracellular Reelin and cytosolic mDab1 (mammalian disabled protein), which activates a tyrosine kinase. This pathway regulates the migration of neurons along the radial glial fiber network during brain

Product Details

	development.
Cross-Reactivity (Details):	Human. Rat.
Purification:	1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target:	VLDLR
Alternative Name:	VLDLR (VLDLR Products)
Background:	Very low-density lipoprotein receptor, VLDL receptor, VLDL-R, VLDLR,VLDL-Receptor (Very Low Density Lipoprotein Receptor) Cellular localisation: Cytoplasmic
Molecular Weight:	143-161kDa
Gene ID:	7436, 370422
UniProt:	P98155
Pathways:	Cellular Response to Molecule of Bacterial Origin

Application Details

Application Notes:	Known_Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT)(Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0 for 45 min at 95°C followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined. Positive_Control: HeLa or U-251 cells. Pancreas, Cerebellum, Heart or Skeletal Muscle.
Restrictions:	For Research Use only

Handling

Concentration:	1.0 mg/mL
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.
Preservative:	Azide free
Storage:	-20 °C,-80 °C
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.
Expiry Date:	24 months