

Datasheet for ABIN6938767

Recombinant anti-VTCN1 antibody



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	VTCN1
Reactivity:	Human
Host:	Rabbit
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This VTCN1 antibody is un-conjugated
Application:	ELISA, Flow Cytometry (FACS), Immunofluorescence (IF), Coating (Coat), Immunohistochemistry (Formalin-fixed Sections) (IHC (f))

Product Details

Immunogen:	A recombinant fragment of human B7-H4 protein (exact sequence is proprietary)
Clone:	B7H4-2652R
Isotype:	IgG
Specificity:	T cell activation and immune function are regulated by the innate immune system through positive and negative costimulatory proteins. One such protein, B7-H4 (B7-homolog 4), belongs to the B7 immunoglobulin superfamily of ligand-lymphocyte interacting proteins. Expressed primarily on the membrane of lymphoid cells, B7-H4 is an immuno-inhibitory protein that interacts with receptors on the surface of T lymphocytes, thus mediating cellular and humoral immune responses. Overexpression of the B7-H4 protein is associated with certain malignancies, including ovarian and breast cancer, as its interaction with T cells suppresses

Product Details

tumor-associated immunity. Current research suggests that, similar to Mucin 16 (CA-125), B7-H4 may be a useful biomarker for the early detection of ovarian cancer.

Cross-Reactivity (Details): Human.

Purification: 1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target: VTCN1

Alternative Name: VTCN1 ([VTCN1 Products](#))

Background: B7 family member, H4, B7 homolog 4, B7 superfamily member 1, B7-H4, B7S1, Immune costimulatory protein B7-H4, T-cell costimulatory molecule B7x, V-set domain-containing T-cell activation inhibitor 1, VCTN1,B7-H4 (Immuno-Inhibitory Protein)
Cellular localisation: Cell Surface & Cytoplasmic

Molecular Weight: 35kDa

Gene ID: 79679, 546434

UniProt: [Q7Z7D3](#)

Application Details

Application Notes: Known_Application: ELISA (Use Ab at 2-4 µg/mL for coating) (Order Ab without BSA), Flow Cytometry (1-2 µg/million cells in 0.1ml), Immunofluorescence (1-2 µg/mL), Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes), Optimal dilution for a specific application should be determined.
Positive_Control: HeLa, SKBr-3 or MCF-7 cells. Human ovary, pancreas, placenta or spleen.

Restrictions: For Research Use only

Handling

Concentration: 1.0 mg/mL

Buffer: Prepared in 10 mM PBS, WITHOUT BSA and Azide.

Preservative: Azide free

Storage: -20 °C,-80 °C

Handling

Storage Comment: Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.

Expiry Date: 24 months