

Datasheet for ABIN6938791

Recombinant anti-YBX1 antibody[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	YBX1
Reactivity:	Human
Host:	Mouse
Antibody Type:	Recombinant Antibody
Clonality:	Monoclonal
Conjugate:	This YBX1 antibody is un-conjugated
Application:	Immunohistochemistry (Formalin-fixed Sections) (IHC (f))

Product Details

Immunogen:	Recombinant human full-length YBX1 protein
Clone:	RYBX1-2430
Isotype:	IgG1 kappa
Specificity:	The specificity of this monoclonal antibody to its intended target was validated by HuProt™ Array, containing more than 19,000, full-length human proteins. Y-box binding protein-1 (YBX1) is the prototypic member of the cold shock protein family that fulfills numerous cellular functions. In the nucleus, YBX1 protein orchestrates transcription of proliferation-related genes, whereas in the cytoplasm it associates with mRNA and directs translation. In human tumor entities, such as breast, lung and prostate cancer, cellular YBX1 expression indicates poor clinical outcome, suggesting that YBX1 is an attractive marker to predict patients' prognosis.
Cross-Reactivity (Details):	Human.

Product Details

Purification: 1.0mg/ml of Ab purified from Bioreactor by Protein A/G.

Target Details

Target:	YBX1
Alternative Name:	YBX1 (YBX1 Products)
Background:	BP-8, CSDA2, CSDB, DBPB, MDR-NF1, NSEP-1, NSEP1, YB-1, YB1,YBX1 / Y-box Binding Protein 1 / YB-1 (Tumor Biomarker) Cellular localisation: Cytoplasmic and Nuclear
Molecular Weight:	36kDa
Gene ID:	4904, 473583
UniProt:	P67809
Pathways:	Regulation of Muscle Cell Differentiation

Application Details

Application Notes:	Known_Application: Immunohistochemistry (Formalin-fixed) (1-2 µg/mL for 30 min at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes), Optimal dilution for a specific application should be determined. Positive_Control: MCF-7, HEK293 or PC12 cells. Prostate, Breast or Lung Carcinoma.
Restrictions:	For Research Use only

Handling

Concentration:	1.0 mg/mL
Buffer:	Prepared in 10 mM PBS, WITHOUT BSA and Azide.
Preservative:	Azide free
Storage:	-20 °C,-80 °C
Storage Comment:	Antibody without azide store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous.
Expiry Date:	24 months