

Datasheet for ABIN6938823

ICOSLG Protein (AA 19-256) (His tag)

Images



	ve	rvi	0	W
\cup	VC	I V I	\sim	v v

Overview	
Quantity:	100 μg
Target:	ICOSLG
Protein Characteristics:	AA 19-256
Origin:	Cynomolgus
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This ICOSLG protein is labelled with His tag.
Product Details	
Purity:	>90 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per μg by the LAL method.
Target Details	
Target:	ICOSLG
Alternative Name:	B7-H2 (ICOSLG Products)
Background:	ICOS ligand (ICOSLG) is also known as B7 homolog 2 (B7-H2), B7-related protein 1 (B7RP-1) and CD antigen CD275, which belongs to the immunoglobulin superfamily and BTN/MOG family. ICOSLG contains one Ig-like C2-type (immunoglobulin-like) domain and one Ig-like V-type (immunoglobulin-like) domain. Isoform 1 is widely expressed, while isoform 2 is detected only in lymph nodes, leukocytes and spleen. B7-H2 is ligand for the T-cell-specific cell surface
	and appearing design and appearing a sign and appearing a sign and appearing design and appearing design and appearing a sign appearing design and appearing a sign appearing a

receptor ICOS. B7-H2 acts as a costimulatory signal for T-cell proliferation and cytokine secretion and induces also B-cell proliferation and differentiation into plasma cells. B7-H2 could play an important role in mediating local tissue responses to inflammatory conditions, as well as in modulating the secondary immune response by co-stimulating memory T-cell function.

Molecular Weight: 28.5 kDa

NCBI Accession: XP_005548617

Pathways: Production of Molecular Mediator of Immune Response, Activated T Cell Proliferation


Application Details

Restrictions: For Research Use only

Handling


Format:	Lyophilized	
Buffer:	PBS, pH 7.4	
Storage:	-20 °C	

Images



ELISA

Image 1. Immobilized Human ICOS, Fc Tag (ABIN6731300,ABIN6809913) at 1 μ g/mL (100 μ L/well) can bind Cynomolgus B7-H2, His Tag (ABIN6933646,ABIN6938823) with a linear range of 0.6-20 ng/mL (QC tested).



SDS-PAGE

 $\label{eq:mage 2.} \textbf{Endown} \textbf{Image 2.} \textbf{ Cynomolgus B7-H2, His Tag on under reducing (R)} \\ \textbf{Condition.} \textbf{ The gel was stained overnight with Coomassie} \\ \textbf{Blue.} \textbf{ The purity of the protein is greater than 90 \%} \ .$